

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Advanced Subsidiary Level and Advanced Level

| CANDIDATE<br>NAME                        |  |                     |                   |  |  |
|--|--|---------------------|-------------------|--|--|
| CENTRE<br>NUMBER                         |  | CANDIDATE<br>NUMBER |                   |  |  |
| CHEMISTRY 9701/22                        |  |                     |                   |  |  |
| Paper 2 Structured Questions AS Core     |  |                     | May/June 2010     |  |  |
|  |  |                     | 1 hour 15 minutes |  |  |
| Candidates answer on the Question Paper. |  |                     |                   |  |  |

Additional Materials: Data Booklet

## READ THESE INSTRUCTIONS FIRST

Write your name, Centre number and candidate number on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs, or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO **NOT** WRITE ON ANY BARCODES.

Answer all questions.

You may lose marks if you do not show your working or if you do not use appropriate units. A Data Booklet is provided.

The number of marks is given in brackets [] at the end of each question or part question. At the end of the examination, fasten all your work securely together.

| For Examiner's Use |  |  |
|--------------------|--|--|
| 1                  |  |  |
| 2                  |  |  |
| 3                  |  |  |
| 4                  |  |  |
| 5                  |  |  |
| Total              |  |  |

This document consists of **12** printed pages.



Answer **all** the questions in the spaces provided.

1 In the 19th and 20th centuries, experimental results showed scientists that atoms consist of a positive, heavy nucleus which is surrounded by electrons.

For Examiner's Use

Then in the 20th century, theoretical scientists explained how electrons are arranged in orbitals around atoms.

- (a) The diagram below represents the energy levels of the orbitals present in atoms of the second period (Li to Ne).
  - (i) Label the energy levels to indicate the principal quantum number **and** the type of orbital at each energy level.



(ii) On the axes below, draw a sketch diagram of **one** of each **different type (shape)** of orbital that is occupied by the electrons in a second-period element.

Label each type.



(iii) Complete the electronic configurations of nitrogen atoms and oxygen atoms on the energy level diagrams below. Use arrows to represent electrons.



2 Copper, proton number 29, and argon, proton number 18, are elements which have different physical and chemical properties.In the solid state, each element has the same face-centred cubic crystal structure which is shown below.



The particles present in such a crystal may be atoms, molecules, anions or cations. In the diagram above, the particles present are represented by .

(a) Which types of particle are present in the copper and argon crystals? In each case, give their formula.

| element | particle | formula |
|---------|----------|---------|
| copper  |          |         |
| argon   |          |         |

[2]

At room temperature, copper is a solid while argon is a gas.

(b) Explain these observations in terms of the forces present in **each** solid structure.



For Examiner's

**3** The table below gives data for some of the oxides of Period 3 elements.

Use oxide Na<sub>2</sub>O MgO  $Al_2O_3$ SiO<sub>2</sub>  $P_4O_6$ SO<sub>2</sub> melting point/°C 24 1275 2827 2017 1607 -75 bonding structure (a) Complete the table by filling in the 'bonding' row by using only the words 'ionic' or 'covalent', (i) (ii) the 'structure' row by using **only** the words 'simple' **or** 'giant'. [2] (b) From the table of oxides above, suggest the formula of one oxide that is completely insoluble in water. [1] ..... (c) Separate samples of Na<sub>2</sub>O and SO<sub>2</sub> were added to water. (i) For **each** oxide, write a balanced equation for its reaction with water and suggest a numerical value for the pH of the resulting solution. Na<sub>2</sub>O equation ..... pH .....  $SO_2$ equation ..... pH ..... Construct a balanced equation for the reaction that occurs when a solution of Na<sub>2</sub>O (ii) in water reacts with a solution of  $SO_2$  in water. ..... [5]

(d) Separate samples of the oxides MgO and SiO<sub>2</sub> are melted.
Each molten sample is then tested to see whether or not it conducts electricity.

Suggest what would be the results in each case. Explain your answers.

[Total: 12]

For

Examiner's Use

- 4 An organic compound, **E**, has the following composition by mass: C, 48.7%; H, 8.1%; O, 43.2%.
  - (a) Calculate the empirical formula of E.

- (b) When vaporised in a suitable apparatus, 0.130 g of E occupied a volume of 58.0 cm<sup>3</sup> at 127 °C and 1.00 × 10<sup>5</sup> Nm<sup>-2</sup>.
  - (i) Use the expression  $pV = \frac{mRT}{M_r}$  to calculate  $M_r$  of **E**, where *m* is the mass of **E**.

(ii) Hence calculate the molecular formula of E.

[4]

For

Examiner's Use

[2]

(c) Compound **F**, is an ester with the molecular formula  $C_4H_8O_2$ .

 ${\bf F}$  is one of four isomers,  ${\bf S},\,{\bf T},\,{\bf U},\,{\rm and}\,\,{\bf V},\,{\rm that}\,\,{\rm are}\,\,{\rm all}\,\,{\rm esters}.$ 

In the boxes below, the structural formula of **S** is given.

Draw the structural formulae of the other **three** isomers of **F** that are esters.

| HCO <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub> |   |   |   |
|--|---|---|---|
| S  | т | U | v |

[3]

| (d) | Wh   | When the ester <b>F</b> is hydrolysed, an alcohol <b>G</b> is produced.  |                   |  |  |
|-----|--|--|-------------------|--|--|
|     | (i)  | What reagent can be used to hydrolyse an ester to an alcohol?  | Examiner's<br>Use |  |  |
|     |  |  |                   |  |  |
|     | (ii)   | What other type of organic compound is produced at the same time?  |                   |  |  |
|     |  | [2]  |                   |  |  |
| (e) | (e) On mild oxidation, the alcohol <b>G</b> gives a compound <b>H</b> which forms a silver mirror with Tollens' reagent. |  |                   |  |  |
|     | (i)  | What functional group does the reaction with Tollens' reagent show to be present in compound $\mathbf{H}$ ? Give the name of this group. |                   |  |  |
|     |  |  |                   |  |  |
|     | (ii)   | What type of alcohol is <b>G</b> ?   |                   |  |  |
|     |  |  |                   |  |  |
|     | (iii)  | What could be the structural formula of the alcohol G?   |                   |  |  |
|     |  |  |                   |  |  |
|     |  | [3]  |                   |  |  |
| (f) | (i)  | Which of the four isomers, <b>S</b> , <b>T</b> , <b>U</b> , or <b>V</b> , could <b>not</b> be <b>F</b> ?                                 |                   |  |  |
|     |  |  |                   |  |  |
|     | (ii)   | Explain your answer.   |                   |  |  |
|     |  |  |                   |  |  |
|     |  | [2]  |                   |  |  |
|     |  | [Total: 16]  |                   |  |  |
|     |  |  |                   |  |  |

**5** Alkenes such as propene can be readily prepared from alcohols in a school or college laboratory by using the apparatus below.

For Examiner's Use



- (ii) Draw the **skeletal** formula of the alcohol you have named in (i).
- (iii) What type of reaction occurs in this case?

.....

[3]

(b) (i) During the reaction, the material X becomes black in colour. Suggest the identity of the black substance and suggest how it is produced during the reaction.

.....

(ii) At the end of the experiment, when no more propene is being produced, the delivery For tube is removed from the water before the apparatus is allowed to cool. Examiner's Use Suggest why this done. ..... ..... (iii) The material labelled **X** can be broken crockery, broken brick or pumice. Give the chemical formula of a compound that is present in one of these materials. ..... (iv) State another reagent that could be used to produce propene from an alcohol. ..... [5] (c) Give the structural formula of the organic product formed when propene reacts separately with each of the following substances. bromine (i) (ii) cold, dilute manganate(VII) ions

11

(iii) hot, concentrated manganate(VII) ions

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.