

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**

Cambridge International Advanced Level

## **MARK SCHEME for the October/November 2015 series**

### **9701 CHEMISTRY**

**9701/42**

Paper 4 (A2 Structured Questions), maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2015 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

bestexamhelp.com

® IGCSE is the registered trademark of Cambridge International Examinations.

<b>Page 2</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International A Level – October/November 2015</b>	<b>9701</b>	<b>42</b>

<b>Question</b>	<b>Marking point</b>	<b>Marks</b>
<b>1 (a)</b>	Ca 3s <sup>2</sup> 3p <sup>6</sup> 4s <sup>2</sup> and Ca <sup>2+</sup> 3s <sup>2</sup> 3p <sup>6</sup>	<b>1</b>
<b>(b)</b>	Ca(OH) <sub>2</sub> + 2HNO <sub>3</sub> → Ca(NO <sub>3</sub> ) <sub>2</sub> + 2H <sub>2</sub> O  or CaO + 2HNO <sub>3</sub> → Ca(NO <sub>3</sub> ) <sub>2</sub> + H <sub>2</sub> O	<b>1</b>
<b>(c) (i)</b>	CaO and brown gas	<b>1</b>
<b>(ii)</b>	the (cat)ion size / radii increases  decreasing its ability to polarise the nitrate ion / N-O bond	<b>2</b>
<b>(d) (i)</b>	(energy change when) 1 mole of ions  <b>gaseous</b> (ions) dissolve in <b>water</b> (to form an infinitely dilute solution) or <b>gaseous</b> (ions) form an <b>aqueous</b> solution	<b>2</b>
<b>(ii)</b>	$\Delta H_{\text{latt}}^{\ominus} \text{Ca(NO}_3)_2 + \Delta H_{\text{sol}}^{\ominus} \text{Ca(NO}_3)_2 = \Delta H_{\text{hyd}}^{\ominus} \text{Ca}^{2+} + 2\Delta H_{\text{hyd}}^{\ominus} \text{NO}_3^{-}$ $\Delta H_{\text{latt}}^{\ominus} - 19 = -1650 + (2x - 314)$  -2259 kJ mol <sup>-1</sup>	<b>3</b>
<b>1</b>	Ca <sup>(2+)</sup> is a smaller (ion) or Ca <sup>(2+)</sup> has a larger charge density Ca <sup>(2+)</sup> has a stronger attraction / bond to H <sub>2</sub> O	<b>2</b>
		<b><u>12</u></b>


<b>Page 3</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International A Level – October/November 2015</b>	<b>9701</b>	<b>42</b>

<b>Question</b>	<b>Marking point</b>	<b>Marks</b>																
<b>2 (a)</b>	<table border="1"> <tr> <td>Na</td> <td>Mg</td> <td>Al</td> <td>Si</td> <td>P</td> <td>S</td> <td>Cl</td> <td>Ar</td> </tr> <tr> <td>1</td> <td>0</td> <td>1</td> <td>2</td> <td>3</td> <td>2</td> <td>1</td> <td>0</td> </tr> </table>	Na	Mg	Al	Si	P	S	Cl	Ar	1	0	1	2	3	2	1	0	<b>3</b>
Na	Mg	Al	Si	P	S	Cl	Ar											
1	0	1	2	3	2	1	0											
<b>(b) (i)</b>	SiCl <sub>4</sub> white solid / ppt <b>or</b> misty / white / steamy fumes pH 0–3 PCl <sub>5</sub> misty / white / steamy fumes pH 0–3	<b>3</b>																
<b>(ii)</b>	SiCl <sub>4</sub> + 2H <sub>2</sub> O → SiO <sub>2</sub> + 4HCl	<b>1</b>																
		<b>7</b>																

<b>Page 4</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International A Level – October/November 2015</b>	<b>9701</b>	<b>42</b>

<b>Question</b>	<b>Marking point</b>	<b>Marks</b>															
<b>3 (a)</b>	forms (one or more) ions with incomplete d orbital(s)/sub-shells/shells	<b>1</b>															
<b>(b) (i)</b>	dative (covalent) <i>or</i> co-ordinate	<b>1</b>															
<b>(ii)</b>	<table border="1"> <thead> <tr> <th>species</th> <th>can act as a ligand</th> <th>cannot act as a ligand</th> </tr> </thead> <tbody> <tr> <td>NO<sub>3</sub><sup>-</sup></td> <td>✓</td> <td></td> </tr> <tr> <td>BF<sub>3</sub></td> <td></td> <td>✓</td> </tr> <tr> <td>H<sub>2</sub>NCH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub></td> <td>✓</td> <td></td> </tr> <tr> <td>NH<sub>4</sub><sup>+</sup></td> <td></td> <td>✓</td> </tr> </tbody> </table>	species	can act as a ligand	cannot act as a ligand	NO <sub>3</sub> <sup>-</sup>	✓		BF <sub>3</sub>		✓	H <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub>	✓		NH <sub>4</sub> <sup>+</sup>		✓	<b>2</b>
species	can act as a ligand	cannot act as a ligand															
NO <sub>3</sub> <sup>-</sup>	✓																
BF <sub>3</sub>		✓															
H <sub>2</sub> NCH <sub>2</sub> CH <sub>2</sub> NH <sub>2</sub>	✓																
NH <sub>4</sub> <sup>+</sup>		✓															
<b>(c) (i)</b>	<table border="1"> <thead> <tr> <th></th> <th><b>formula of manganese species formed</b></th> <th><b>type of reaction</b></th> </tr> </thead> <tbody> <tr> <td>Mn<sup>2+</sup> (aq) + NaOH (aq)</td> <td>Mn(OH)<sub>2</sub> Mn(H<sub>2</sub>O)<sub>4</sub>(OH)<sub>2</sub> Mn(OH)<sub>3</sub></td> <td>precipitation</td> </tr> <tr> <td>Mn<sup>2+</sup> (aq) + concentrated HCl</td> <td>MnCl<sub>4</sub><sup>2-</sup> MnCl<sub>6</sub><sup>4-</sup></td> <td>ligand exchange / substitution</td> </tr> <tr> <td>Mn<sup>2+</sup> (aq) + aqueous H<sub>2</sub>O<sub>2</sub></td> <td>Mn<sup>3+</sup></td> <td>redox / oxidation</td> </tr> </tbody> </table>		<b>formula of manganese species formed</b>	<b>type of reaction</b>	Mn <sup>2+</sup> (aq) + NaOH (aq)	Mn(OH) <sub>2</sub> Mn(H <sub>2</sub> O) <sub>4</sub> (OH) <sub>2</sub> Mn(OH) <sub>3</sub>	precipitation	Mn <sup>2+</sup> (aq) + concentrated HCl	MnCl <sub>4</sub> <sup>2-</sup> MnCl <sub>6</sub> <sup>4-</sup>	ligand exchange / substitution	Mn <sup>2+</sup> (aq) + aqueous H <sub>2</sub> O <sub>2</sub>	Mn <sup>3+</sup>	redox / oxidation	<b>5</b>			
	<b>formula of manganese species formed</b>	<b>type of reaction</b>															
Mn <sup>2+</sup> (aq) + NaOH (aq)	Mn(OH) <sub>2</sub> Mn(H <sub>2</sub> O) <sub>4</sub> (OH) <sub>2</sub> Mn(OH) <sub>3</sub>	precipitation															
Mn <sup>2+</sup> (aq) + concentrated HCl	MnCl <sub>4</sub> <sup>2-</sup> MnCl <sub>6</sub> <sup>4-</sup>	ligand exchange / substitution															
Mn <sup>2+</sup> (aq) + aqueous H <sub>2</sub> O <sub>2</sub>	Mn <sup>3+</sup>	redox / oxidation															
		<b>9</b>															

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2015	9701	42

Question	Marking point	Marks
4 (a)	<p>M1: dipole on C–Cl bond</p> <p>M2: curly arrow breaking C–Cl bond</p> <p>M3: curly arrow from the oxygen on <math>\text{OH}^-</math> (lone pair needs to be shown) to carbon in C–Cl bond <b>and</b> <math>\text{Cl}^-</math> (ion) formed in the mechanism</p> 	3
(b) (i)	time taken for the concentration of a reactant(s) to fall to half its original value	1
(ii)	evidence of a pair of construction lines on graph <b>and</b> $t_{1/2} = 49\text{--}53$ s	1
(iii)	no effect/change	1
(c) (i)	evidence of tangent at 80 s <b>and</b> data used, e.g. $0.42/152 = 0.00263$ units $\text{mol dm}^{-3}\text{s}^{-1}$	2
(ii)	correct use of answer to (i)/0.19 <b>and</b> $\text{s}^{-1}$	1
		<u>9</u>


<b>Page 6</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International A Level – October/November 2015</b>	<b>9701</b>	<b>42</b>

<b>Question</b>	<b>Marking point</b>	<b>Marks</b>
<b>5 (a) (i)</b>	M1: salt bridge <b>and</b> voltmeter/ M2: method of H <sub>2</sub> gas delivery M3: X <b>and</b> Pt electrode labelled M4: solution H <sup>+</sup> /HCl(aq)/H <sub>2</sub> SO <sub>4</sub> <b>and</b> X <sup>2+</sup> labelled	<b>4</b>
<b>(ii)</b>	25 °C/298 K <b>and</b> 1 atm/101 kPa pressure <b>and</b> 1 mol dm <sup>-3</sup> (solution)	<b>1</b>
<b>(iii)</b>	solution – ions <b>or</b> H <sup>+</sup> and X <sup>2+</sup> <b>and</b> wires – electrons/e <sup>-</sup>	<b>1</b>
<b>(b) (i)</b>	$X + 2Ag^+ \rightarrow 2Ag + X^{2+}$	<b>1</b>
<b>(ii)</b>	moles Ag = 1.30 / 107.9 = 0.0120 1 moles of X react with 2 moles Ag <sup>+</sup> moles of X lost = 0.012 × 0.5 = 0.00602 A <sub>r</sub> of X = 0.67/0.006 = 111–112 <b>and</b> X = Cd	<b>4</b>
		<b><u>11</u></b>


Page 7	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2015	9701	42

Question	Marking point	Marks
6 (a)	$4\text{BF}_3 + 3\text{NaBH}_4 \rightarrow 2\text{B}_2\text{H}_6 + 3\text{NaBF}_4$	1
(b)	<p>[1] dipoles (M1)</p> <p>[1] intermediate (M3)</p> <p>[1] both curly arrows (M2) arrow <u>must</u> come from lone pair</p>	3
(c) (i)	(electrophilic) addition	1
(ii)		1

Page 8	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2015	9701	42

Question	Marking point	Marks
(d) (i)	<p>any four of</p> <p>M1: <math>\sigma</math>-bonds between C–C or C–H</p> <p>M2: <math>\pi</math>-bonds formed from overlap of p-orbitals</p> <p>M3: (<math>\pi</math>-bonds/electrons) above and below the ring</p> <p>M4: bonds/electrons are delocalised</p> <p>M5: bond angle <math>120^\circ</math></p> <p>M6: intermediate C–C bond length / all C–C same length / strength</p> <p>M7: carbons are <math>sp^2</math> hybridised</p>	3
(ii)	<p>correct delocalised structure of borazine</p> 	1
		<b>10</b>

Page 9	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2015	9701	42

Question	Marking point	Marks
7 (a) (i)		3
(ii)	<p><math>\text{Sn} + \text{HCl}</math></p> <p><math>\text{HNO}_2</math> or <math>\text{NaNO}_2 + \text{HCl}</math></p> <p>step 1 (linked to a reduction) reflux/heat/<math>&gt;50^\circ\text{C}</math> <b>or</b> conc/6M (HCl)  <b>and</b> step 2 <math>\leq 10^\circ\text{C}</math></p>	3
(iii)	diazonium (group)	1
(b) (i)	<p><math>\sigma</math>-bonds = 14</p> <p><math>\pi</math>-bonds = 2</p>	2


Page 10	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2015	9701	42

Question	Marking point	Marks									
7	<table border="1"> <thead> <tr> <th>reagent</th> <th>structure of product</th> <th>type of reaction</th> </tr> </thead> <tbody> <tr> <td>HCl</td> <td> </td> <td>acid-base or neutralisation</td> </tr> <tr> <td>CH<sub>3</sub>CH<sub>2</sub>Br</td> <td> </td> <td>(nucleophilic) substitution</td> </tr> </tbody> </table>	reagent	structure of product	type of reaction	HCl		acid-base or neutralisation	CH <sub>3</sub> CH <sub>2</sub> Br		(nucleophilic) substitution	3
reagent	structure of product	type of reaction									
HCl		acid-base or neutralisation									
CH <sub>3</sub> CH <sub>2</sub> Br		(nucleophilic) substitution									
		<u>12</u>									



<b>Page 11</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International A Level – October/November 2015</b>	<b>9701</b>	<b>42</b>

<b>Question</b>	<b>Marking point</b>	<b>Marks</b>
<b>8 (a) (i)</b>	<b>A</b> = mRNA <b>B</b> <sub>1</sub> and <b>B</b> <sub>2</sub> , etc. = tRNA <b>or</b> tRNA-amino acid complex	<b>2</b>
<b>(ii)</b>	stage 1 = transcription <b>and</b> stage 3= translation	<b>1</b>
<b>(b) (i)</b>	C <sub>5</sub> H <sub>5</sub> N <sub>5</sub>	<b>1</b>
<b>(ii)</b>	cytosine, thymine, guanine	<b>1</b>
<b>(iii)</b>	covalent hydrogen bonding	<b>2</b>
<b>(c)</b>	hydrolysis	<b>1</b>
<b>(d) (i)</b>	Phosphorus / P	<b>1</b>
<b>(ii)</b>	H atoms have insufficient electron density <b>or</b> electrons (to show up) <b>or</b> H atoms contain one e <sup>-</sup>	<b>1</b>
		<b><u>10</u></b>

Page 12	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2015	9701	42

Question	Marking point	Marks
9 (a)	iron/Fe (= haemoglobin) sodium/Na or potassium/K (= transmission of nerve impulses) Zn or Cu or Mg or Mn or Mo or Ni or Fe or Co (= enzyme co-factor)	2
(b)	any three of: M1: substrate binds to/fits into the <b>active site</b> of the enzyme M2: Interaction with site causes a specific bond to be weakened, (which breaks) M3: lowers activation energy M4: products released from the enzyme/active site	3
(c) (i)	Tertiary	1
(ii)	$2 -SH \rightarrow -S - S- (+ 2H)$	1
(iii)	oxidation	1
(d) (i)	<b>E = CH and F = CH<sub>2</sub></b>	1
(ii)	<b>E = triplet and adjacent 2H</b> <b>F = doublet and adjacent 1H</b>	2
		<u>11</u>
10 (a) (i)		1

Page 13	Mark Scheme	Syllabus	Paper
	Cambridge International A Level – October/November 2015	9701	42

Question	Marking point	Marks
(ii)	 <p>The diagram shows a chemical structure of a dipeptide derivative. It consists of a benzyl group attached to a chiral carbon, which is also bonded to a methoxy ester group (-OCH<sub>3</sub>) and a hydrogen atom. This chiral carbon is linked to an amide group (-NH-), which is further bonded to another chiral carbon. This second chiral carbon is bonded to an amino group (-NH<sub>2</sub>) and a carboxylic acid group (-COOH). Two arrows point to the oxygen atom of the ester group and the hydrogen atom of the amide group, respectively.</p>	2
(iii)	 <p>The diagram shows three chemical structures. On the left is a dipeptide derivative with a benzyl group, a carboxylic acid group, and an amino group. In the middle is the structure of glycine (2-aminoacetic acid). On the right is the structure of methanol (CH<sub>3</sub>-OH).</p>	3
(b)	<p>M1: hydrogen bonding  M2: between the NH<sub>2</sub> groups and water  or CO<sub>2</sub>/C=O/-OH groups and water (allow names)  or lone pair on N/O with water</p>	2
(c)	allow range 1–200 nm or 1–200 × 10 <sup>-9</sup> m	1
		<u>9</u>