Cambridge International **AS & A Level** 

896665

∞

# Cambridge International Examinations

Cambridge International Advanced Subsidiary and Advanced Level

CANDIDATE NAME					
CENTRE NUMBER		CANDIDATE NUMBER			
CHEMISTRY 9701/22					
Paper 2 AS Le	vel Structured Questions	February/March 2016			
		1 hour 15 minutes			

Candidates answer on the Question Paper.

Additional Materials: Data Booklet

#### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen. You may use an HB pencil for any diagrams or graphs. Do not use staples, paper clips, glue or correction fluid. DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions. Electronic calculators may be used. You may lose marks if you do not show your working or if you do not use appropriate units. A Data Booklet is provided.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

This document consists of 13 printed pages and 3 blank pages.



2

Answer **all** the questions in the spaces provided.

- 1 This question is about Period 3 elements and their compounds.
  - (a) Give an explanation for each of the following statements.
    - (i) The atomic radius decreases across Period 3 (Na to Ar).

(iv) Magnesium is a better electrical conductor than sodium.

.....[1]

- 3
- (b) The flow chart below shows a series of reactions.



2 Spathose is an iron ore that contains iron(II) carbonate, FeCO<sub>3</sub>. The percentage of iron(II) carbonate in spathose can be determined by titration with acidified potassium dichromate(VI) solution using a suitable indicator.

The ionic equation is shown below.

 $Cr_2O_7^{2-}(aq) + 14H^+(aq) + 6Fe^{2+}(aq) \rightarrow 2Cr^{3+}(aq) + 6Fe^{3+}(aq) + 7H_2O(I)$ 

(a) A 5.00 g sample of spathose was reacted with excess concentrated hydrochloric acid and then filtered.

The filtrate was made up to 250 cm<sup>3</sup> in a volumetric flask with distilled water.

A 25.0 cm<sup>3</sup> sample of the standard solution required 27.30 cm<sup>3</sup> of 0.0200 mol dm<sup>-3</sup> dichromate(VI) solution for complete reaction.

(i) Calculate the amount, in moles, of dichromate(VI) ions used in the titration.

amount = ..... mol [1]

(ii) Use your answer to (i) to calculate the amount, in moles, of Fe<sup>2+</sup> present in the 25.0 cm<sup>3</sup> sample.

amount = ..... mol [1]

(iii) Use your answer to (ii) to calculate the amount, in moles, of Fe<sup>2+</sup> present in the 250 cm<sup>3</sup> volumetric flask.

amount = ..... mol [1]

(iv) Use your answer to (iii) to calculate the mass of iron(II) carbonate present in the sample of spathose.

mass = ..... g [2]

(v) Calculate the percentage of iron(II) carbonate in the sample of spathose.

percentage of iron(II) carbonate = ...... % [1]

(b) Iron ores containing iron(III) compounds can be analysed using a similar method.

A standard solution of an aqueous iron(III) compound is reacted with aqueous tin(II) chloride. Aqueous tin(IV) chloride and aqueous iron(II) chloride are the products of this reaction.

(i) Write an ionic equation for this reaction. Do not include state symbols.

(ii) Any excess tin(II) chloride can be removed by reaction with HgC $l_2(aq)$ . A white precipitate of Hg<sub>2</sub>C $l_2$  is produced.

Complete the equation for this reaction.

$$\dots(\dots) + \dots \operatorname{HgC} l_2(\operatorname{aq}) \to \operatorname{SnC} l_4(\dots) + \operatorname{Hg}_2 \operatorname{C} l_2(\dots)$$

[2]

[Total: 10]

**3** Over one million tonnes of hydrogen cyanide, HCN, are produced each year using the Andrussow process. The overall equation for the reaction is shown.

 $CH_4(g) + NH_3(g) + 1\frac{1}{2}O_2(g) \rightleftharpoons HCN(g) + 3H_2O(g)$ 

(a) (i) Draw a dot-and-cross diagram to represent the bonding in a molecule of ammonia, NH<sub>3</sub>, and state the shape of the molecule.



The bonding between the carbon and nitrogen atoms consists of one sigma ( $\sigma$ ) bond and two pi ( $\pi$ ) bonds.

Sketch the shape of the sigma bond and one of the pi bonds in the space below. Show clearly the position of the atomic nuclei in each diagram.



- (b) The reaction exists as a dynamic equilibrium.
  - (i) Explain what is meant by the term *dynamic equilibrium*.

.....

.....[1]

(ii) State and explain how the amounts of the chemicals present in the equilibrium mixture will change when the pressure is increased.

(c) The process uses a platinum catalyst, which increases the rate of reaction.

Sketch a Boltzmann distribution on the axes given below and use your diagram to explain how the platinum catalyst increases the rate of the reaction.



- (d) The reaction of hydrogen cyanide with propanone is an important first step in many organic syntheses.
  - (i) Give the full name of the mechanism of this reaction.

(ii) Complete the diagram to show the mechanism of the reaction of hydrogen cyanide with propanone.

Draw the structure of the intermediate and the product of the reaction. Include all relevant charges, partial charges, curly arrows and lone pairs.



[Total: 17]

Question 4 starts on the next page.

4 The following compounds were all found to be components of a sample of petrol.

$$\begin{array}{cccc} & H & CH_{3} \\ CH_{3}(CH_{2})_{2}CH_{3} & (CH_{3})_{3}CCH_{2}CH(CH_{3})_{2} & H_{3}C & -C & -C & -OH \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ \end{array}$$
(i) Give the molecular formula of compound **G**.
  
(ii) Give the empirical formula of compound **H**.
  
[1]

- ......[1]
- (iii) Draw the **skeletal** formula of compound **J**.

(b)	) Write an equation to represent the complete combustion of compound <b>H</b> .				
	[1]				
(C)	Fossil fuels are often contaminated with sulfur.				
	State and explain why supplies of fossil fuels that contain sulfur pose a problem to the environment.				

(a)

(d) The boiling points of compounds G, H and J are shown below.

compound	G	Н	J
boiling point/°C	0	99	112

Explain the differences in the boiling points of the three compounds.

[4]

(e) Compound J can be produced from 2-chloro-3-methylbutane,  $C_5H_{11}Cl$ .

Give the reagent(s) and conditions for this reaction.

......[1]

[Total: 11]

**5** Some reactions of compound **P**,  $C_5H_8O$ , are shown.



(a) (i) Give the structures for organic compounds Q, R, S and T.





wavenumber/cm<sup>-1</sup>

Use the information given to suggest a structure for compound  ${\bf U}.$  Explain your answer.

## **BLANK PAGE**

## **BLANK PAGE**

#### **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.