

Cambridge International Examinations

Cambridge International Advanced Subsidiary and Advanced Level

CHEMISTRY		9701/34
CENTRE NUMBER	CANDIDATE NUMBER	
CANDIDATE NAME		

Paper 3 Advanced Practical Skills 2

May/June 2016

2 hours

Candidates answer on the Question Paper.

As listed in the Confidential Instructions Additional Materials:

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Give details of the practical session and laboratory where appropriate, in the boxes provided.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 10 and 11.

A copy of the Periodic Table is printed on page 12.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

Session
Laboratory

For Exam	iner's Use
1	
2	
3	
Total	

This document consists of 11 printed pages and 1 blank page.



1 Borax is an alkali which has many uses. In this experiment you will determine **x** in the chemical formula of borax, Na₂B_xO₇.10H₂O, by titration with hydrochloric acid.

FB 1 is a solution containing 15.5 g dm⁻³ of borax, $Na_2B_xO_7$.10 H_2O . **FB 2** is 2.00 mol dm⁻³ hydrochloric acid, HC *l*. methyl orange indicator

(a) Method

Dilution of FB 2

- Pipette 10.0 cm³ of FB 2 into the 250 cm³ volumetric flask.
- Make the solution up to 250 cm³ using distilled water.
- Shake the solution in the volumetric flask thoroughly.
- This diluted solution of hydrochloric acid is **FB 3**. Label the volumetric flask **FB 3**.

Titration

- Fill the burette with **FB 3**.
- Pipette **25.0 cm³** of **FB 1** into a conical flask.
- Add several drops of methyl orange.
- Perform a rough titration and record your burette readings in the space below.

The rough	titre is		cm ³
-----------	----------	--	-----------------

[7]

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make sure any recorded results show the precision of your practical work.
- Record in a suitable form below all of your burette readings and the volume of FB 3 added in each accurate titration.

I II III IV V VI VII

(b) From your accurate titration results, obtain a suitable value for the volume of **FB 3** to be used in your calculations.

Show clearly how you obtained this value.

25.0 cm³ of **FB 1** required cm³ of **FB 3**. [1]

(c) Calculation	าร
-----------------	----

Show your working and appropriate significant figures	in the final answer to each step of your
calculations.	

cal	culations.
(i)	Calculate the number of moles of hydrochloric acid present in the volume of FB 3 calculated in (b) .
	(2).
	moles of $HCl = \dots mol$
(ii)	1 mole of borax is neutralised by 2 moles of hydrochloric acid. Calculate the number of moles of borax that react with the hydrochloric acid in (i).
	moles of borax = mol
(iii)	Use your answer to (ii) to calculate the number of moles of borax in 1.00 dm³ of FB 1.
	moles of borax in 1.00 dm ³ FB 1 = mol
(iv)	Use your answer to (iii) and the information on page 2 to calculate the relative formula mass, M_r , of borax.
	$M_{\rm r}$ of borax =

(v) Calculate $\bf x$ in the formula of borax, Na₂B_xO₇.10H₂O. Use data from the Periodic Table on page 12.

x = [5]

[Total: 13]

2 Some metal carbonates cannot be obtained in a pure state. For example magnesium carbonate exists in a 'basic' form, in which magnesium hydroxide is also present.

One possible chemical formula of basic magnesium carbonate is MgCO₃.Mg(OH)₂.2H₂O.

When basic magnesium carbonate is heated, if the possible formula were correct, it would decompose as shown below.

$$MgCO_3.Mg(OH)_2.2H_2O(s) \rightarrow 2MgO(s) + CO_2(g) + 3H_2O(g)$$

In this experiment, you will decompose basic magnesium carbonate by heating it, and you will use your results to determine whether this possible formula is correct.

FB 4 is basic magnesium carbonate.

(a) Method

Read through the method before starting any practical work and prepare a table for your results in the space below.

- Weigh a crucible with its lid and record the mass.
- Add 1.1-1.3g of FB 4 to the crucible. Weigh the crucible and lid with FB 4 and record the
 mass.
- Place the crucible on the pipe-clay triangle and remove the lid.
- Heat the crucible and contents **gently** for about one minute.
- Then heat the crucible and contents strongly for about four minutes.
- Replace the lid and allow the crucible to cool for at least five minutes.
- While the crucible is cooling, you may wish to begin work on Question 3.
- Re-weigh the crucible and contents with lid. Record the mass.
- Calculate, and record, the mass of **FB 4** used and the mass of residue obtained.

I II III IV V

[5]

((b)) Calculations
۸	~	, oaioaiatioiio

Show your working and appropriate significant figures in the final answer to each ste	ер	of y	our
calculations.			

(i)	Use your results to calculate the number of moles of magnesium oxide, MgO, obtained as residue.
(ii)	moles of MgO obtained =
	M_{r} of basic magnesium carbonate (from experiment) =
(iii)	Use data from the Periodic Table to calculate the relative formula mass, M_r , of basic magnesium carbonate from its possible formula, ${\rm MgCO_3.Mg(OH)_2.2H_2O}$.
	$M_{\rm r}$ of basic magnesium carbonate (from formula) =
(iv)	If the relative formula mass of basic magnesium carbonate obtained from your experiment is within 2.5% of the answer in (iii), this is good evidence that the possible formula, MgCO ₃ .Mg(OH) ₂ .2H ₂ O, is correct. Does your experiment support the possible formula? Give a reason for your answer.
	[5]

(c)	Eval	luation

(i)	State one way in which the accuracy of the experimental procedure could have been improved using the same mass of FB 4 . Explain your answer.
(ii)	A student carried out the experiment twice using different masses of FB 4 . He used the mean mass of FB 4 and the mean mass of magnesium oxide obtained to calculate the relative formula mass of basic magnesium carbonate.
	Instead of doing this, he could have calculated the relative formula mass of basic magnesium carbonate from his two experiments separately.
	Suggest one advantage of carrying out separate calculations for each experiment.
(iii)	State the error when making one reading on your balance.
	error = g
	Calculate the maximum percentage error in the mass of FB 4 used.
	percentage error = % [4]
	[Total: 14]

BLANK PAGE

3 Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations**.

You should indicate clearly at what stage in a test a change occurs.

No additional tests for ions present should be attempted.

If any solution is warmed, a boiling tube MUST be used.

Rinse and reuse test-tubes and boiling tubes where possible.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

(a) FB 5, FB 6 and FB 7 are solutions, each of which contain one cation and one anion. The anions present are all listed on page 11.

Use a 1 cm depth of these solutions in a test-tube for each of the following tests. Complete the table below.

40.04		observations				
test	FB 5	FB 6	FB 7			
Add a 2 cm strip of magnesium ribbon.						
Add aqueous sodium hydroxide.						
Add an equal depth of aqueous potassium iodide.						
Add a few drops of FB 5 .						

(b) (i)	From the observation made when potassium iodide was added to FB 6 , suggest the identity of the cation in FB 6 . Explain your conclusion.
	cation in FB 6
	explanation
(ii)	FB 5 gives no precipitate when aqueous ammonia is added. Suggest the identities of both ions in FB 5 .
	cation in FB 5
	anion in FB 5
(iii)	Identify FB 7.
(iv)	Give the ionic equation for the reaction between magnesium and FB 7 .
	[4]
	8 is a solid. Carry out the following tests on FB 8. cord your observations in each test.
(i)	Heat a small spatula measure of FB 8 gently in a hard-glass test-tube.
	observations
(ii)	To a 1 cm depth of hydrochloric acid in a test-tube, add a small spatula measure of FB 8 .
	observations
(iii)	What conclusions, if any, can you make about the identities of the ions in FB 8 ?
	cation in FB 8
	anion in FB 8
	[4]

[Total: 13]

Qualitative Analysis Notes

Key: [ppt. = precipitate]

1 Reactions of aqueous cations

	reaction with								
ion	NaOH(aq)	NH ₃ (aq)							
aluminium, A <i>l</i> ³+(aq)	white ppt. soluble in excess	white ppt. insoluble in excess							
ammonium, NH ₄ +(aq)	no ppt. ammonia produced on heating	_							
barium, Ba²+(aq)	faint white ppt. is nearly always observed unless reagents are pure	no ppt.							
calcium, Ca ²⁺ (aq)	white ppt. with high [Ca ²⁺ (aq)]	no ppt.							
chromium(III), Cr³+(aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess							
copper(II), Cu ²⁺ (aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution							
iron(II), Fe ²⁺ (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess							
iron(III), Fe³+(aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess							
magnesium, Mg²+(aq)	white ppt. insoluble in excess	white ppt. insoluble in excess							
manganese(II), Mn²+(aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess							
zinc, Zn ²⁺ (aq)	white ppt. soluble in excess	white ppt. soluble in excess							

2 Reactions of anions

ion	reaction
carbonate, CO ₃ ²⁻	CO ₂ liberated by dilute acids
chloride, C <i>l</i> ⁻ (aq)	gives white ppt. with Ag+(aq) (soluble in NH ₃ (aq))
bromide, Br ⁻ (aq)	gives cream ppt. with Ag ⁺ (aq) (partially soluble in NH ₃ (aq))
iodide, I ⁻ (aq)	gives yellow ppt. with Ag ⁺ (aq) (insoluble in NH ₃ (aq))
nitrate, NO ₃ ⁻ (aq)	NH ₃ liberated on heating with OH ⁻ (aq) and A <i>l</i> foil
nitrite, NO ₂ ⁻ (aq)	NH_3 liberated on heating with $OH^-(aq)$ and Al foil; NO liberated by dilute acids (colourless $NO \rightarrow$ (pale) brown NO_2 in air)
sulfate, SO ₄ ²⁻ (aq)	gives white ppt. with Ba2+(aq) (insoluble in excess dilute strong acids)
sulfite, SO ₃ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (soluble in excess dilute strong acids)

3 Tests for gases

gas	test and test result
ammonia, NH ₃	turns damp red litmus paper blue
carbon dioxide, CO ₂	gives a white ppt. with limewater (ppt. dissolves with excess CO ₂)
chlorine, Cl ₂	bleaches damp litmus paper
hydrogen, H ₂	"pops" with a lighted splint
oxygen, O ₂	relights a glowing splint

The Periodic Table of Elements

	~		ന	Ę C		ന	Ē (N	T	_	<u>ج</u> ه		_	u. 8		(I)	ς ε.		_	5						
	18	2	Ĭ	heliu 4.0	10	ž	neo 20.:	18	₹	argon 39.9	36	Ž	krypt 83.8	75	×	xenc 131	86	ž	radc						
	17				6	Щ	fluorine 19.0	17	Cl	chlorine 35.5	32	Ŗ	bromine 79.9	53	П	iodine 126.9	82	Ąŧ	astatine -						
	16				8	0	oxygen 16.0	16	ഗ	sulfur 32.1	34	Se	selenium 79.0	52	Ц	tellurium 127.6	84	Ро	polonium –	116	_	livermorium	ı		
	15				7	z	nitrogen 14.0	15	۵	phosphorus 31.0	33	As	arsenic 74.9	51	Sp	antimony 121.8	83	<u>.</u>	bismuth 209.0						
	14				9	O	carbon 12.0	14	:S	silicon 28.1	32	Ge	germanium 72.6	20	S	tin 118.7	82	Pp	lead 207.2	114	Εl	flerovium	1		
	13				2	В	boron 10.8	13	Ρl	aluminium 27.0	31	Ga	gallium 69.7	49	In	indium 114.8	81	11	thallium 204.4						
										12	30	Zu	zinc 65.4	48	р О	cadmium 112.4	80	БĤ	mercury 200.6	112	ပ်	copernicium	-		
										1	29	Cu	copper 63.5	47	Ag	silver 107.9	62	Αn	gold 197.0	111	Rg	roentgenium	-		
dn										10	28	Z	nickel 58.7	46	Pd	palladium 106.4	78	₫	platinum 195.1	110	S	darmstadtium	-		
Group										6	27	ပိ	cobalt 58.9	45	R	rhodium 102.9	77	'n	iridium 192.2	109	¥	meitnerium	-		
		-	I	hydrogen 1.0						80	26	Ь	iron 55.8	44	Ru	ruthenium 101.1	92	SO	osmium 190.2	108	¥	hassium	1		
				J					7	25	M	manganese 54.9	43	ပ	technetium -	75	Re	rhenium 186.2	107	В	pohrium				
						00	SS			9	24	ပ်	chromium 52.0	42	Mo	molybdenum 95.9	74	>	tungsten 183.8	106	Sg	seaborgium	1		
						Key	atomic number	atomic symbo	name relative atomic mass			2	23	>	vanadium 50.9	41	g	niobium 92.9	73	Та	tantalum 180.9	105	9	dubnium	-
					al	ator	relat			4	22	F	titanium 47.9	40	Zr	zirconium 91.2	72	Έ	hafnium 178.5	104	꿒	rutherfordium			
										က	21	Sc	scandium 45.0	39	>	yttrium 88.9	57-71	lanthanoids		89–103	actinoids				
	2				4	Be	beryllium 9.0	12	Mg	magnesium 24.3	20	Ca	calcium 40.1	88	ഗ്	strontium 87.6	26	Ba	barium 137.3	88	Ra	radium			
	_				8	:=	lithium 6.9	11	Na	sodium 23.0	19	×	potassium 39.1	37	Rb	rubidium 85.5	55	S	caesium 132.9	87	占	francium			

71	P	lutetium 175.0	103	۲	lawrencium	ı	
		ytterbium 173.1					
69	T	thulium 168.9	101	Md	mendelevium	ı	
89	Ē	erbium 167.3	100	Fm	ferminm	ı	
29	웃	holmium 164.9	66	Es	einsteinium	ı	
99	Δ	dysprosium 162.5	86	Ç	californium	ı	
99	Тр	terbium 158.9	26	益	berkelium	ı	
64	Вd	gadolinium 157.3	96	Cm	curium	ı	
63	Еп	europium 152.0	92	Am	americium	ı	
62	Sm	samarium 150.4	94	Pu	plutonium	ı	
61	Pm	promethium —	93	ď	neptunium	ı	
09	PZ	neodymium 144.4	92	\supset	uranium	238.0	
59	Ā	praseodymium 140.9	91	Ра	protactinium	231.0	
58	Ce	cerium 140.1	06	T	thorium	232.0	
22	Гa	lanthanum 138.9	88	Ac	actinium	ı	

lanthanoids

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.