



Cambridge International AS & A Level

CHEMISTRY

9701/22

Paper 2 AS Level Structured Questions

May/June 2020

MARK SCHEME

Maximum Mark: 60

Published

Students did not sit exam papers in the June 2020 series due to the Covid-19 global pandemic.

This mark scheme is published to support teachers and students and should be read together with the question paper. It shows the requirements of the exam. The answer column of the mark scheme shows the proposed basis on which Examiners would award marks for this exam. Where appropriate, this column also provides the most likely acceptable alternative responses expected from students. Examiners usually review the mark scheme after they have seen student responses and update the mark scheme if appropriate. In the June series, Examiners were unable to consider the acceptability of alternative responses, as there were no student responses to consider.

Mark schemes should usually be read together with the Principal Examiner Report for Teachers. However, because students did not sit exam papers, there is no Principal Examiner Report for Teachers for the June 2020 series.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the June 2020 series for most Cambridge IGCSE™ and Cambridge International A & AS Level components, and some Cambridge O Level components.

This document consists of **12** printed pages.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Science-Specific Marking Principles

1	Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.
2	The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.
3	Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).
4	The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.
5	<p><u>'List rule' guidance</u> (see examples below)</p> <p>For questions that require <i>n</i> responses (e.g. State two reasons ...):</p> <ul style="list-style-type: none">• The response should be read as continuous prose, even when numbered answer spaces are provided• Any response marked <i>ignore</i> in the mark scheme should not count towards <i>n</i>• Incorrect responses should not be awarded credit but will still count towards <i>n</i>• Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should not be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response• Non-contradictory responses after the first <i>n</i> responses may be ignored even if they include incorrect science.

6 Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.


For answers given in standard form, (e.g. $a \times 10^n$) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

7 Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.


Question	Answer	Marks								
1(a)	<p>EITHER</p> <p>M1 (weighted) average/mean mass of the isotope(s)/an atom(s)</p> <p>M2 relative to 1/12 of the mass (of an atom) of ^{12}C (where an atom of ^{12}C is exactly 12).</p> <p>OR</p> <p>M1 mass of one mol of atoms</p> <p>M2 relative / compared to 1/12 (the mass) of 1 mol of C-12 OR in which one mol C-12 (atom) has a mass of (exactly) 12 g</p>	2								
1(b)	<p>M1 $60.11/100 \times 69 + 39.89/100 \times 71$</p> <p>M2 69.80</p>	2								
1(c)	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 15%;">isotope</th> <th style="width: 15%;">nucleon number</th> <th style="width: 30%;">total number of electrons in lowest energy level</th> <th style="width: 40%;">type of orbital contains the electron in the highest energy level</th> </tr> </thead> <tbody> <tr> <td style="text-align: center;">^{71}Ga</td> <td style="text-align: center;">M1 71</td> <td style="text-align: center;">M2 2</td> <td style="text-align: center;">M3 p (-orbital)</td> </tr> </tbody> </table>	isotope	nucleon number	total number of electrons in lowest energy level	type of orbital contains the electron in the highest energy level	^{71}Ga	M1 71	M2 2	M3 p (-orbital)	3
isotope	nucleon number	total number of electrons in lowest energy level	type of orbital contains the electron in the highest energy level							
^{71}Ga	M1 71	M2 2	M3 p (-orbital)							
1(d)	<p>M1 shape</p> <div style="text-align: center;">  </div> <p>M2 bond angle 120°</p>	2								
1(e)(i)	$\text{Ga}_2\text{O}_3 + 6\text{HCl} \rightarrow 2\text{GaCl}_3 + 3\text{H}_2\text{O}$	1								

Question	Answer	Marks
1(e)(ii)	<p>M1 Identity of correct gallium containing product NaGa(OH)₄ OR NaGaO₂</p> <p>M2 correctly balanced equation for reaction of Ga₂O₃ with NaOH(aq) EITHER Ga₂O₃+ 2NaOH + 3H₂O → 2NaGa(OH)₄ OR Ga₂O₃+ 2NaOH → 2NaGaO₂ + H₂O</p>	2


Question	Answer	Marks
2(a)	4NH ₃ + 5O ₂ → 4NO + 6H ₂ O	1
2(b)	$\begin{array}{c} \text{xx} \\ \text{H} \overset{\circ}{\text{N}} \overset{\circ}{\text{H}} \\ \text{x} \quad \text{x} \\ \text{xO} \\ \text{H} \end{array}$	1
2(c)(i)	<p>3NO₂ + H₂O → 2HNO₃ + NO</p> <p>(+4) (+)5 (+)2</p> <p>3 correct oxidation numbers – 2 marks 2 correct oxidation numbers – 1 mark</p>	2
2(c)(ii)	<p>relates the term disproportionation to the reaction described</p> <p>M1 <u>nitrogen /N</u> (in nitrogen dioxide) is both gaining electrons and losing electrons during the reaction</p> <p>M2 refer to relevant transfer of electrons when NO₂ reacts to form HNO₃ and NO</p> <p>NO₂ to HNO₃ involves loss of electron(s)</p> <p>AND</p> <p>NO₂ to NO involves gain of electron(s)</p>	


Question	Answer	Marks
2(d)	M1 state the effect of NO gas on contact with moist air NO reacts with water OR NO reacts with oxygen and water. M2 consequence of M1 in terms of atmospheric pollution causing acid rain OR photochemical smog / ground level ozone OR destroy ozone layer	2
2(e)	M1 number of mol in sample of NH_4NO_3 40t NH_4NO_3 = 500 000 mol ammonium nitrate M2 ratio of mol NO_2 : NH_4NO_3 3/2 mol NO_2 :1 mol NH_4NO_3 M3 volume of no mol NO_2 at rtp using 1 mol = 24dm^3 18 000 000 dm^3 of NO_2	3
2(f)	fertiliser	1


Question	Answer	Marks
3(a)	hydrolysis	1
3(b)	M1 both have molecular formula - $\text{C}_6\text{H}_{12}\text{O}_6$ M2 idea that in glucose and fructose there are the same number and type of atoms present but the atoms are arranged in a different order ie one has a carbonyl group at the end of the chain/molecule and the other has a carbonyl group in the middle of the chain/molecule	2
3(c)(i)	value for the activation energy for the reaction A (no enzyme) compared to reaction B (with enzyme) value / range of values that are more (+) 29 kJ mol^{-1}	1
3(c)(ii)	value for the enthalpy change for reaction A (no enzyme) compared to reaction B (with enzyme) -14 kJ mol^{-1}	1

Question	Answer	Marks
3(c)(iii)	<p>M1 show the energy of the reactants > products AND label ΔH using the predicted value given in (ii)</p> <p>M2 show activation energy 'hump' AND label using the value given in (i)</p> 	2
3(d)(i)	<p>M1 (enthalpy change) when 1 mole of sucrose</p> <p>M2 burns/combusts/reacts in excess air/oxygen</p> <p>OR</p> <p>completely burns/combusts/reacts in air/oxygen</p>	2
3(d)(ii)	<p>M1 for finding amount of energy released per gram of sucrose using $\Delta H/Q = mc\Delta T$ OR $\Delta H = -mc\Delta T$ $= (-)250 \times 4.18 \times (40.7 - 25) = (-)16406.5$ J per gram OR $(-)16.4065$ kJ / g</p> <p>M2 for finding amount (mol) sucrose in 1g = $1/342$ mol</p> <p>M3 = $M1 / (M2 \times 1000)$ $\Delta H = -5610$ kJ mol⁻¹ (3 sig figs) OR -5611 kJ mol⁻¹ (4 sig figs)</p>	3

Question	Answer	Marks
4(a)(i)	Bromine / Br	1
4(a)(ii)	$\text{Ag}^+(\text{aq}) + \text{X}^-(\text{aq}) \rightarrow \text{AgX}(\text{s})$ OR $\text{Ag}^+(\text{aq}) + \text{Br}^-(\text{aq}) \rightarrow \text{AgBr}(\text{s})$	1
4(a)(iii)	<p>M1 reagent Add (aqueous) ammonia</p> <p>M2 expected result EITHER (Dilute ammonia) – partial amount precipitate dissolves OR not much precipitate dissolves OR add concentrated ammonia – precipitate dissolves</p>	2

Question	Answer	Marks
4(b)(i)	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{Br}$	1
4(b)(i)	M1 primary/ 1° (carbo)cation formed is not very stable M2 EITHER (as) only one alkyl group exerting an inductive effect OR only one alkyl group so the charge is (more) localised on the C^+	2
4(c)(i)	elimination	1
4(c)(ii)	$\text{C}_4\text{H}_9\text{Cl} + \text{NaOH} \rightarrow \text{C}_4\text{H}_8 + \text{NaCl} + \text{H}_2\text{O}$	1
4(c)(iii)		1
4(c)(iv)	M1 2-chloro(-2-)methylpropane M2 1-chloro(-2-)methylpropane ALLOW in any order	2

Question	Answer			Marks
5(a)	Rxn.	name of mechanism	Name of reagents and conditions	6
1	M1 electrophilic addition	M2 steam AND concentrated phosphoric acid (catalyst)		
2		M3 & M4 Two marks for name of reagent and both conditions. One mark for name of reagent and one conditions acidified potassium dichromate ((VI)) condition 1 warm condition 2 distil NOT reflux		
6	M5 nucleophilic substitution	M6 ammonia (alcoholic) AND heat in a sealed tube / heat under pressure		
5(b)(i)	<p><i>mechanism for ethanal and HCN using CN⁻ (from KCN) as the catalyst</i></p>  <p>M1 arrow from lone pair of electrons on :CN⁻ to C of C=O M2 correct dipole on C^{δ+}=O^{δ-} AND arrow from the double bond to or beyond the O of C=O M3 arrow from lone pair of electrons on O of intermediate to H of HCN AND arrow from H-C bond to C of H-C≡N</p>			3

Question	Answer	Marks
5(b)(ii)	ALLOW in any order M1 nitrile M2 (secondary) alcohol	2
5(b)(iii)	 <p>ALLOW any unambiguous structure</p>	1