

Surname	Centre Number	Candidate Number
Other Names		2



GCE A LEVEL

1410U50-1E



CHEMISTRY – A2 unit 5 Practical Methods and Analysis Task

FRIDAY, 10 MAY 2019 – MORNING

1 hour

For Examiner's use only		
Question	Maximum Mark	Mark Awarded
1.	13	
2.	8	
3.	9	
Total	30	

1410U501E
01

ADDITIONAL MATERIALS

In addition to this examination paper, you will need a:

- calculator, pencil and ruler;
- **Data Booklet** supplied by WJEC.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen.

Write your name, centre number and candidate number in the spaces at the top of this page.

Answer **all** questions in the spaces provided.


INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

The maximum mark for this paper is 30.

Your answers must be relevant and must make full use of the information given to be awarded full marks for a question.

If you run out of space, use the additional page(s) at the back of the booklet, taking care to number the question(s) correctly.

- 
- methyl benzenecarboxylate + $\text{HNO}_3(\text{conc}) \xrightarrow{\text{H}_2\text{SO}_4(\text{conc})}$ compound **A** + H_2O

Stage 1

Stage 2

Stage 3

Stage 4

Once addition was complete, the reaction mixture was kept at room temperature for a further 15 minutes before it was poured onto a small amount of crushed ice in a beaker. Solid compound **A** was formed.

- (a) Suggest why the reaction mixture was cooled during stage 3.

[1]

.....

.....

- (b) Compound **A** is very much more soluble in hot ethanol than it is in cold ethanol.

Describe how you would purify the sample of compound **A**.

[3]

.....

.....

.....

.....

.....

.....

- (c) After purification 4.56 g of compound **A** was isolated.


Calculate the percentage yield of this reaction.

[4]


(density of methyl benzenecarboxylate is 1.08 g cm^{-3})

Percentage yield = %


- (d) $C_8H_7NO_4$ has a number of different isomers. Three of them are shown below.



compound **A**



5-aminobenzene-1,3-dicarboxylic acid



1-(4-hydroxy-3-nitrophenyl)ethanone

- (i) State **two** chemical tests that will give a positive result for 5-aminobenzene-1,3-dicarboxylic acid but **not** for compound **A**.

Give the reagent(s) and observation(s) for a positive test.

[2]

Test 1

Reagent(s)

Observation(s)

Test 2

Reagent(s)

Observation(s)

- (ii) State **three** chemical tests that will give a positive result for 1-(4-hydroxy-3-nitrophenyl)ethanone but **not** for compound **A**.

Give the reagent(s) and observation(s) for a positive test.

[3]

Test 1

Reagent(s)

Observation(s)

Test 2

Reagent(s)

Observation(s)

Test 3

Reagent(s)

Observation(s)

1410U501E
05

13

CO_3^{2-}	I^-	Cl_2	$\text{S}_2\text{O}_3^{2-}$
carbonate	iodide	chlorine	thiosulfate

- dilute H_2SO_4
- $\text{AgNO}_3(\text{aq})$

You should include observations and **ionic** equations for any reactions occurring. [8]

BLANK PAGE

3. A solid mixture contains sodium carbonate, sodium hydrogencarbonate and an unreactive impurity. It is known to contain 62.3 % by mass of sodium carbonate.

A student prepared a solution of the solid mixture by dissolving 8.72 g of it in water, transferring the solution into a 500 cm³ volumetric flask and making up to the mark with deionised water. The solution was shaken vigorously and labelled as solution **X**.

The percentage by mass of sodium hydrogencarbonate in the solid mixture was then determined by two different methods.

Method 1: Titration


- 25.0 cm³ of solution **X** was transferred into a conical flask and 3-5 drops of phenolphthalein indicator added.
- At the phenolphthalein end-point (indicator colour change pink to colourless) the following reaction is complete.



- 0.196 mol dm⁻³ hydrochloric acid was added from a burette whilst swirling, until the first permanent colour change occurred. The volume of hydrochloric acid used (volume **V**₁) was recorded.
- 3-5 drops of methyl orange indicator were added to the solution in the conical flask.
- At the methyl orange end-point (indicator colour change yellow to pink) the following reaction is complete.



- The titration was continued with more of the hydrochloric acid added from the burette whilst swirling, until the first permanent colour change occurred. The **additional volume** of hydrochloric acid used (volume **V**₂) was recorded.
- After carrying out one rough titration, the titration was repeated several times and mean values for **V**₁ and **V**₂ calculated.

Summary**Results**

- Mean volume V_1 of hydrochloric acid used = 13.10 cm^3
- Mean volume V_2 of hydrochloric acid used = 20.80 cm^3

(a) Why did the student repeat the titration?

[1]

.....

.....

(b) (i) Explain why the difference between volume V_2 and V_1 represents the volume of acid required to react with the original sodium hydrogencarbonate in the solid mixture.

[1]

.....

.....

.....

.....

(ii) Use the difference between volume V_2 and V_1 to calculate the percentage by mass of sodium hydrogencarbonate in the original solid mixture.

[3]

Percentage by mass = %

Method 2: Gas collection

- 25.0 cm³ of solution **X** was transferred to a conical flask and excess hydrochloric acid added.
- 99.7 cm³ of carbon dioxide gas was formed and collected in a gas syringe.

(Assume the reaction was carried out at 298 K and 1 atm pressure)

- (c) (i) The original solid mixture contained 62.3 % by mass of sodium carbonate. Of the 99.7 cm³ of carbon dioxide gas produced in this reaction, calculate the volume produced by the reaction between hydrochloric acid and the sodium carbonate in 25.0 cm³ of solution **X**. [2]

Volume = cm³

- (ii) Hence, calculate the percentage by mass of sodium hydrogencarbonate in the original solid mixture. [2]

Percentage by mass = %

END OF PAPER

