
MATHEMATICS

9709/13

Paper 1

October/November 2016

MARK SCHEME

Maximum Mark: 75

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2016 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

| | | | |
|--------|--|----------|-------|
| Page 2 | Mark Scheme | Syllabus | Paper |
| | Cambridge International AS/A Level – October/November 2016 | 9709 | 13 |

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol ∇ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
 - Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

| | | | |
|---------------|---|-----------------|--------------|
| Page 3 | Mark Scheme | Syllabus | Paper |
| | Cambridge International AS/A Level – October/November 2016 | 9709 | 13 |

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF/OE Any Equivalent Form (of answer is equally acceptable) / Or Equivalent

AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)

CAO Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)

CWO Correct Working Only – often written by a ‘fortuitous’ answer

ISW Ignore Subsequent Working

SOI Seen or implied

SR Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through $\frac{1}{2}$ ” marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR –2 penalty may be applied in particular cases if agreed at the coordination meeting.

PA –1 This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

| Page 4 | Mark Scheme | Syllabus | Paper |
|--------|--|----------|-------|
| | Cambridge International AS/A Level – October/November 2016 | 9709 | 13 |

| | | | | |
|-------|---|---|-----|--|
| 1 | $kx^2 - 3x = x - k \Rightarrow kx^2 - 4x + k (= 0)$ $(-4)^2 - 4(k)(k)$ soi $k > 2$, $k < -2$ cao Allow $(2, \infty)$ etc. Allow $2 < k < -k$ | M1 M1 A1 | [3] | Eliminate y and rearrange into 3-term quad $b^2 - 4ac$. |
| 2 | $(+/-)20 \times 3^3(x^3)$, $10a^3(x^3)$ soi $-540 + 10a^3 = 100$ oe $a = 4$ | B1B1 M1 A1 | [4] | Each term can include x^3 Must have 3 terms and include a^3 and 100 |
| 3 | $4\sin^2 x = 6\cos^2 x \Rightarrow \tan^2 x = \frac{6}{4}$ or $4\sin^2 x = 6(1 - \sin^2 x)$ $[\tan x = (\pm)1.225$ or $\sin x = (\pm)0.7746$ or $\cos x = (\pm)0.6325]$ $x = 50.8$ (Allow 0.886 (rad)) Another angle correct $x = 50.8^\circ, 129.2^\circ, 230.8^\circ, 309.2^\circ$ [0.886, 2.25/6, 4.03, 5.40 (rad)] | M1 A1 A1 ^h A1 | [4] | Or $4(1 - \cos^2 x) = 6\cos^2 x$ Or any other angle correct Ft from 1st angle (Allow radians) All 4 angles correct in degrees |
| 4 | $f'(x) = 3x^2 - 6x - 9$ soi Attempt to solve $f'(x) = 0$ or $f'(x) > 0$ or $f'(x) \geq 0$ soi $(3)(x-3)(x+1)$ or 3, -1 seen or 3 only seen Least possible value of n is 3. Accept $n = 3$. Accept $n \geq 3$ | B1 M1 A1 A1 | [4] | With or without equality/inequality signs Must be in terms of n |
| 5 (i) | $\cos 0.9 = OE / 6$ or $= \sin\left(\frac{\pi}{2} - 0.9\right)$ oe $OE = 6\cos 0.9 = 3.73$ oe AG | M1 A1 | [2] | Other methods possible |
| (ii) | Use of $(2\pi - 1.8)$ or equivalent method Area of large sector $= \frac{1}{2} \times 6^2 \times (2\pi - 1.8)$ oe Area of small sector $\frac{1}{2} \times 3.73^2 \times 1.8$ Total area $= 80.7(0) + 12.5(2) = 93.2$ | M1 M1 M1 A1 | [4] | Expect 4.48 Or $\pi 6^2 - \frac{1}{2} 6^2 1.8$. Expect 80.70 Expect 12.52 Other methods possible |
| 6 (i) | $\frac{2+x}{2} = n \Rightarrow x = 2n - 2$ $\frac{m+y}{2} = -6 \Rightarrow y = -12 - m$ | B1 B1 | [2] | No MR for $(\frac{1}{2}(2+n), \frac{1}{2}(m-6))$ Expect $(2n-2, -12-m)$ |

| Page 5 | Mark Scheme | Syllabus | Paper |
|--------|--|----------|-------|
| | Cambridge International AS/A Level – October/November 2016 | 9709 | 13 |

| | | | | |
|-------|---|---|-----|---|
| (ii) | Sub <i>their</i> x, y into $y = x + 1 \rightarrow -12 - m = 2n - 2 + 1$ $\frac{m+6}{2-n} = -1$ oe Not nested in an equation Eliminate a variable $m = -9, n = -1$ | M1* B1 DM1 A1A1 | [5] | Expect $m + 2n = -11$ Expect $m - n = -8$ Note: other methods possible |
| 7 (i) | AB.AC = $3 - 2 - 1 = 0$ hence perpendicular or 90° AB.AD = $3 + 4 - 7 = 0$ hence perpendicular or 90° AC.AD = $1 - 8 + 7 = 0$ hence perpendicular or 90° AG | B1 B1 B1 | [3] | $3 - 2 - 1$ or sum of prods etc must be seen Or single statement: mutually perpendicular or 90° seen at least once . |
| (ii) | Area $ABC = (\frac{1}{2})\sqrt{3^2 + 1^2 + 1^2} \times \sqrt{1^2 + (-2)^2 + (-1)^2}$ $= \frac{1}{2}\sqrt{11} \times \sqrt{6}$ Vol. = $\frac{1}{3} \times \text{their } \Delta ABC \times \sqrt{1^2 + 4^2 + (-7)^2}$ $= \frac{1}{6}\sqrt{66} \times \sqrt{66} = 11$ | M1 A1 M1 A1 | [4] | Expect $\frac{1}{2}\sqrt{66}$ Not 11.0 |
| 8 (i) | $(2x+3)^2 + 1$ Cannot score retrospectively in (iii) | B1B1B1 | [3] | For $a = 2, b = 3, c = 1$ |
| (ii) | $g(x) = 2x + 3$ cao | B1 | [1] | In (ii),(iii) Allow if from $4\left(x + \frac{3}{2}\right)^2 + 1$ |
| (iii) | $y = (2x+3)^2 + 1 \Rightarrow 2x+3 = (\pm)\sqrt{y-1}$ or ft from (i) $x = (\pm)\frac{1}{2}\sqrt{y-1} - \frac{3}{2}$ or ft from (i) $(fg)^{-1}(x) = \frac{1}{2}\sqrt{x-1} - \frac{3}{2}$ cao Note alt. method $g^{-1}f^{-1}$ Domain is $(x) > 10$ ALT. method for first 3 marks: Trying to obtain $g^{-1}[f^{-1}(x)]$ $g^{-1} = \frac{1}{2}(x-3), f^{-1} = \sqrt{x-1}$ A1 for $\frac{1}{2}\sqrt{x-1} - \frac{3}{2}$ | M1 M1 A1 B1 *M1 DM1 A1 | [4] | Or with x/y transposed. Or with x/y transposed Allow sign errors. Must be a function of x . Allow $y = \dots$ Allow $(10, \infty), 10 < x < \infty$ etc. but not with y or f or g involved. Not ≥ 10 Both required |

| Page 6 | Mark Scheme | Syllabus | Paper |
|--------|--|----------|-------|
| | Cambridge International AS/A Level – October/November 2016 | 9709 | 13 |

| | | | | |
|--------|--|--|-----|---|
| 9 (a) | $\frac{6}{1-r} = \frac{12}{1+r}$ $r = \frac{1}{3}$ $S = 9$ | M1 A1 A1 | [3] | |
| (b) | $\frac{13}{2}[2\cos\theta + 12\sin^2\theta] = 52$ $2\cos\theta + 12(1 - \cos^2\theta) = 8 \rightarrow 6\cos^2\theta - \cos\theta - 2 (= 0)$ $\cos\theta = 2/3 \text{ or } -1/2 \text{ soi}$ $\theta = 0.841, 2.09 \text{ Dep on previous A1}$ | M1* DM1 A1 A1A1 | [5] | Use of correct formula for sum of AP Use $s^2 = 1 - c^2$ & simplify to 3-term quad Accept $0.268\pi, 2\pi/3$. SRA1 for $48.2^\circ, 120^\circ$ Extra solutions in range -1 |
| 10 (i) | at $x = a^2, \frac{dy}{dx} = \frac{2}{a^2} + \frac{1}{a^2}$ or $2a^{-2} + a^{-2} \left(= \frac{3}{a^2} \text{ or } 3a^{-2} \right)$ $y - 3 = \frac{3}{a^2}(x - a^2) \text{ or } y = \frac{3}{a^2}x + c \rightarrow 3 = \frac{3}{a^2}a^2 + c$ $y = \frac{3}{a^2}x \text{ or } 3a^{-2}x \text{ cao}$ | B1 M1 A1 | [3] | $\frac{2}{a^2} + \frac{1}{a^2}$ or $2a^{-2} + a^{-2}$ seen anywhere in (i) Through $(a^2, 3)$ & with <i>their</i> grad as $f(a)$ |
| (ii) | $(y) = \frac{2}{a} x^{1/2} + \frac{ax^{-1/2}}{-1/2} (+c)$ sub $x = a^2, y = 3$ into $\int dy/dx$ $c = 1 \left(y = \frac{4x^{1/2}}{a} - 2ax^{-1/2} + 1 \right)$ | B1B1 M1 A1 | [4] | c must be present. Expect $3 = 4 - 2 + c$ |
| (iii) | sub $x = 16, y = 8 \rightarrow 8 = \frac{4}{a} \times 4 - 2a \times \frac{1}{4} + 1$ $a^2 + 14a - 32 (= 0)$ $a = 2$ $A = (4, 3), B = (16, 8) \quad AB^2 = 12^2 + 5^2 \rightarrow AB = 13$ | *M1 A1 A1 DM1A1 | [5] | Sub into <i>their</i> y Allow -16 in addition |

| Page 7 | Mark Scheme | Syllabus | Paper |
|--------|--|----------|-------|
| | Cambridge International AS/A Level – October/November 2016 | 9709 | 13 |

| | | | | |
|----------------------|---|---|------------|--|
| <p>11 (i)</p> | <p>Attempt diffn. and equate to 0 $\frac{dy}{dx} = -k(kx-3)^{-2} + k = 0$</p> <p>$(kx-3)^2 = 1$ or $k^3x^2 - 6k^2x + 8k (= 0)$</p> <p>$x = \frac{2}{k}$ or $\frac{4}{k}$</p> <p>$\frac{d^2y}{dx^2} = 2k^2(kx-3)^{-3}$</p> <p>When $x = \frac{2}{k}$, $\frac{d^2y}{dx^2} = (-2k^2) < 0$ MAX All previous</p> <p>When $x = \frac{4}{k}$, $\frac{d^2y}{dx^2} = (2k^2) > 0$ MIN working correct</p> | <p>*M1</p> <p>DM1</p> <p>*A1*A1</p> <p>B1⁴</p> <p>DB1</p> <p>DB1</p> | <p>[7]</p> | <p>Must contain $(kx-3)^{-2}$ + other term(s)</p> <p>Simplify to a quadratic</p> <p>Legitimately obtained</p> <p>It must contain $Ak^2(kx-3)^{-3}$ where $A > 0$</p> <p>Convincing alt. methods (values either side) must show which values used & cannot use $x = 3/k$</p> |
| <p>(ii)</p> | <p>$V = (\pi) \int [(x-3)^{-1} + (x-3)]^2 dx$</p> <p>$= (\pi) \int [(x-3)^{-2} + (x-3)^2 + 2] dx$</p> <p>$= (\pi) \left[-(x-3)^{-1} + \frac{(x-3)^3}{3} + 2x \right]$ Condone missing 2x</p> <p>$= (\pi) \left[1 - \frac{1}{3} + 4 - \left(\frac{1}{3} - 9 + 0 \right) \right]$</p> <p>$= 40\pi / 3$ oe or 41.9</p> | <p>*M1</p> <p>A1</p> <p>A1</p> <p>DM1</p> <p>A1</p> | <p>[5]</p> | <p>Attempt to expand y^2 and then integrate</p> <p>Or</p> <p>$\left[-(x-3)^{-1} + \frac{x^3}{3} - 3x^2 + 9x + 2x \right]$</p> <p>Apply limits $0 \rightarrow 2$</p> <p>2 missing $\rightarrow 28\pi / 3$ scores M1A0A1M1A0</p> |