

Cambridge International Examinations Cambridge International Advanced Level

* 7 2 6 4 7 0 3 4 6 7 *	CANDIDATE NAME		
	CENTRE NUMBER	CANDIDATE NUMBER	
	MATHEMATICS	3	9709/71
	Paper 7 Probab	bility & Statistics 2 (S2)	May/June 2017
			1 hour 15 minutes
	Candidates ans	wer on the Question Paper.	
	Additional Mater	rials: List of Formulae (MF9)	
	READ THESE I	NSTRUCTIONS FIRST	

Write your Centre number, candidate number and name in the spaces at the top of this page. Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 50.

This document consists of 11 printed pages and 1 blank page.



- 1 On average, 1 clover plant in 10 000 has four leaves instead of three.
 - (i) Use an approximating distribution to calculate the probability that, in a random sample of 2000 clover plants, more than 2 will have four leaves. [3]

..... (ii) Justify your approximating distribution. [2]

2 Past experience has shown that the heights of a certain variety of plant have mean 64.0 cm and standard deviation 3.8 cm. During a particularly hot summer, it was expected that the heights of plants of this variety would be less than usual. In order to test whether this was the case, a botanist recorded the heights of a random sample of 100 plants and found that the value of the sample mean was 63.3 cm. Stating a necessary assumption, carry out the test at the 2.5% significance level. [6]

 •••••
 •••••
 •••••

(a) The waiting time at a certain bus stop has variance 2.6 minutes². For a random sample of 3 75 people, the mean waiting time was 7.1 minutes. Calculate a 92% confidence interval for the population mean waiting time. [3]

(b) A researcher used 3 random samples to calculate 3 independent 92% confidence intervals. Find the probability that all 3 of these confidence intervals contain only values that are greater than the actual population mean. [2] (c) Another researcher surveyed the first 75 people who waited at a bus stop on a Monday morning. Give a reason why this sample is unsuitable for use in finding a confidence interval for the mean waiting time. [1]



6

The time, X minutes, taken by a large number of runners to complete a certain race has probability density function f given by

$$f(x) = \begin{cases} \frac{k}{x^2} & 5 \le x \le 10, \\ 0 & \text{otherwise,} \end{cases}$$

where *k* is a constant, as shown in the diagram.

(i) Without calculation, explain how you can tell that there were more runners whose times were below 7.5 minutes than above 7.5 minutes. [1]



(iii)	Find E(<i>X</i>). [3]
		••••
		•••
(iv)	Find Var(X).	2]
(iv)	Find Var(X).	2]
(iv)	Find Var(<i>X</i>).	2]
(iv)	Find Var(X).	2]
(iv)		····
(iv)		····
(iv)		····
(iv)		···· ····
(iv)		···· ····
(iv)		···· ····
(iv)		

5 Large packets of sugar are packed in cartons, each containing 12 packets. The weights of these packets are normally distributed with mean 505 g and standard deviation 3.2 g. The weights of the cartons, when empty, are independently normally distributed with mean 150 g and standard deviation 7 g.

(i)	Find the probability that the total weight of a full carton is less than 6200 g. [5]	5]
		•••
		•••
		•••
		•••
		•••
		•••
		•••
		•••
		•••
		•••
		•••
		•••
		•••
		•••
		•••

Small packets of sugar are packed in boxes. The total weight of a full box has a normal distribution with mean 3130 g and standard deviation 12.1 g.

(ii) Find the probability that the weight of a randomly chosen full carton is less than double the weight of a randomly chosen full box. [5]

.....

6 The number of sports injuries per month at a certain college has a Poisson distribution. In the past the mean has been 1.1 injuries per month. The principal recently introduced new safety guidelines and she decides to test, at the 2% significance level, whether the mean number of sports injuries has been reduced. She notes the number of sports injuries during a 6-month period.

(i)	Find the critical region for the test and state the probability of a Type I error. [6]
(ii)	State what is meant by a Type I error in this context. [1]

(iii)	During the 6-month period there are a total of 2 sports injuries. Carry out the test.	[3]
(iv)	Assuming that the mean remains 1.1, calculate the probability that there will be fewer than	30
		[4]
		•••••

BLANK PAGE

12

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.