

Cambridge International Examinations

Cambridge International Advanced Level

CANDIDATE NAME			
CENTRE NUMBER		CANDIDATE NUMBER	
MATHEMATICS			9709/52
Paper 5 Mechan	nics 2 (M2)	October/Novem	nber 2018
		1 hour 15	5 minutes
Candidates answ	wer on the Question Paper.		
Additional Mater	ials: List of Formulae (MF9)		

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

Where a numerical value for the acceleration due to gravity is needed, use 10 m s^{-2} .

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 50.



International Examinations

	ontal ground. Find the speed of B when the path of B makes an angle of 20° above the horizontal ground.
•••••	
•••••	
•••••	
•••••	
base	iform solid object is made by attaching a cone to a cylinder so that the circumferences of the cone and a plane face of the cylinder coincide. The cone and the cylinder each have read and height 0.4 m.
base 0.3 m	of the cone and a plane face of the cylinder coincide. The cone and the cylinder each have r
base 0.3 m (i)	of the cone and a plane face of the cylinder coincide. The cone and the cylinder each have read and height 0.4 m. Calculate the distance of the centre of mass of the object from the vertex of the cone.
base 0.3 m (i)	of the cone and a plane face of the cylinder coincide. The cone and the cylinder each have r and height $0.4 \mathrm{m}$.
base 0.3 m (i)	of the cone and a plane face of the cylinder coincide. The cone and the cylinder each have read and height 0.4 m. Calculate the distance of the centre of mass of the object from the vertex of the cone.
base 0.3 m (i)	of the cone and a plane face of the cylinder coincide. The cone and the cylinder each have reach height 0.4 m. Calculate the distance of the centre of mass of the object from the vertex of the cone. [The volume of a cone is $\frac{1}{3}\pi r^2h$.]
base 0.3 m (i)	of the cone and a plane face of the cylinder coincide. The cone and the cylinder each have read and height 0.4 m. Calculate the distance of the centre of mass of the object from the vertex of the cone. [The volume of a cone is $\frac{1}{3}\pi r^2h$.]
base 0.3 m (i)	of the cone and a plane face of the cylinder coincide. The cone and the cylinder each have reach height 0.4 m. Calculate the distance of the centre of mass of the object from the vertex of the cone. [The volume of a cone is $\frac{1}{3}\pi r^2h$.]
base 0.3 m (i)	of the cone and a plane face of the cylinder coincide. The cone and the cylinder each have reach height 0.4 m. Calculate the distance of the centre of mass of the object from the vertex of the cone. [The volume of a cone is $\frac{1}{3}\pi r^2h$.]
base 0.3 m (i)	of the cone and a plane face of the cylinder coincide. The cone and the cylinder each have reach height 0.4 m. Calculate the distance of the centre of mass of the object from the vertex of the cone. [The volume of a cone is $\frac{1}{3}\pi r^2h$.]

						•••••
						••••••
	•••••	•••••			•••••	• • • • • • • • • • • • • • • • • • • •
					•••••	•••••
						•••••
f magnitude <i>kW</i> does not slip.	N acting at 30)° to the upwa	ard vertical is a	pplied to the v	vertex of the	cone. T
of magnitude <i>kW</i> does not slip.	N acting at 30)° to the upwa	ard vertical is a	pplied to the v	vertex of the	cone. Ti
oject has weight Vof magnitude kW does not slip.	N acting at 30)° to the upwa	ard vertical is a	pplied to the v	vertex of the	cone. Tl
of magnitude kW does not slip.	N acting at 30)° to the upwa	ard vertical is a	pplied to the v	vertex of the	surface. cone. Tl
f magnitude <i>kW</i> does not slip.	N acting at 30)° to the upwa	ard vertical is a	pplied to the v	vertex of the	cone. Ti
f magnitude <i>kW</i> does not slip.	N acting at 30)° to the upwa	ard vertical is a	pplied to the v	vertex of the	cone. T
magnitude <i>kW</i> loes not slip.	N acting at 30)° to the upwa	ard vertical is a	pplied to the v	vertex of the	cone. T
magnitude <i>kW</i> loes not slip.	N acting at 30)° to the upwa	ard vertical is a	pplied to the v	vertex of the	cone. T
f magnitude <i>kW</i> does not slip.	N acting at 30)° to the upwa	ard vertical is a	pplied to the v	vertex of the	cone. T
f magnitude <i>kW</i> does not slip.	N acting at 30)° to the upwa	ard vertical is a	pplied to the v	vertex of the	cone. T
of magnitude <i>kW</i> does not slip.	N acting at 30)° to the upwa	ard vertical is a	pplied to the v	vertex of the	cone. Ti
of magnitude <i>kW</i> does not slip.	N acting at 30)° to the upwa	ard vertical is a	pplied to the v	vertex of the	cone. T
f magnitude <i>kW</i> does not slip.	N acting at 30)° to the upwa	ard vertical is a	pplied to the v	vertex of the	cone. Ti

3	Afte 8x N One	article P of mass 0.4 kg is projected horizontally along a smooth horizontal plane from a point O . er projection the velocity of P is v m s ⁻¹ and its displacement from O is x m. A force of magnitude V directed away from V 0 acts on V 1 and a force of magnitude (V 2e ^{-x} + 4) V 1 opposes the motion of V 2 end of a light elastic string of natural length 0.5 m is attached to V 2 and the other end of the string stached to V 3.
	(i)	Show that $v \frac{dv}{dx} = 20x - 10 - 5e^{-x}$ before the elastic string becomes taut. [2]
	(ii)	Given that the initial velocity of P is $6 \mathrm{ms^{-1}}$, find v when the string first becomes taut. [3]

When the string is taut, the acceleration of P is proportional to e^{-x} .

(iii)	Find the modulus of elasticity of the string.	[2]
		· • • • • •
		· • • • • •
		· • • • • •
		· • • • • •
		· • • • • •
		.
		•••••
		· • • • • •
		· • • • • •
		· • • • • •
		.
		· • • • • •
		· • • • • •


	nd y m respectiv	ely.			
) Exp	ress x and y in	terms of t and t	nence show that the	e equation of the p	ath of the object is
v =	$-\frac{5x^2}{V^2}.$				[3]
,	V^2				[-]
•••••		•••••		•••••	
•••••					
• • • • •		•••••	••••••	•••••	
• • • • • •					
• • • • •		•••••		•••••	••••••
•••••					
••••					
	ct passes throu	gh points with co	pordinates $(a, -a)$	and $(a^2, -16a)$, where	here a is a positive
obje	et passes amou				
	er passes anou,				•
tant.					
tant.	If the value of a .				
ant.					
tant.					[3]
stant.					
stant.					
stant.					

(iii)	Given that the object strikes the ground at the point where $x = 5a$, find the height of Q above the
(iii)	Given that the object strikes the ground at the point where $x = 5a$, find the height of O above the ground.
(iii)	Given that the object strikes the ground at the point where $x = 5a$, find the height of O above the ground.
(iii)	
(iii)	
(iii)	
(iii)	ground . [2]

A particle P of mass $0.7 \, \mathrm{kg}$ is attached to a fixed point O by a light elastic string of natural length

Find th	ne distance	below A of	the point a	nt which P	comes to i	nstantaneou	s rest.	
•••••		••••••	••••••		••••••	•••••••	••••••	•••••
						•••••		•••••
								•••••
						•••••		•••••
•••••		•••••	••••••		••••••	•••••	•••••	•••••
		•••••				•••••		
						• • • • • • • • • • • • • • • • • • • •		
•••••								•••••
						•••••		
		•••••	••••••		••••••	•••••	•••••	•••••
						•••••		
		•••••	••••••		••••••	•••••	•••••	•••••
		•••••				•••••		
	•••••		•••••			• • • • • • • • • • • • • • • • • • • •	•••••	•••••


Find the greatest speed of <i>P</i> in the motion.	
	•••••



The diagram shows a uniform lamina ABCDEFGH. The lamina consists of a quarter-circle OAB of radius r m, a rectangle DEFG and two isosceles right-angled triangles COD and GOH. The rectangle has DG = EF = r m and DE = FG = x m.

given that the centre of i	mass of the lamina is at O , express x in te	rms of r .

Civan instead that the materials DEEC is a square with adopt of langth up, state with a	
Given instead that the rectangle $DEFG$ is a square with edges of length r m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle	
Given instead that the rectangle $DEFG$ is a square with edges of length r m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	ason
Given instead that the rectangle $DEFG$ is a square with edges of length r m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle $DEFG$ is a square with edges of length r m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle $DEFG$ is a square with edges of length r m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle $DEFG$ is a square with edges of length r m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle $DEFG$ is a square with edges of length r m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle $DEFG$ is a square with edges of length r m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle $DEFG$ is a square with edges of length r m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle $DEFG$ is a square with edges of length r m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle $DEFG$ is a square with edges of length r m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle <i>DEFG</i> is a square with edges of length <i>r</i> m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle $DEFG$ is a square with edges of length r m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle <i>DEFG</i> is a square with edges of length <i>r</i> m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle <i>DEFG</i> is a square with edges of length <i>r</i> m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle <i>DEFG</i> is a square with edges of length <i>r</i> m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle <i>DEFG</i> is a square with edges of length <i>r</i> m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle <i>DEFG</i> is a square with edges of length <i>r</i> m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	
Given instead that the rectangle <i>DEFG</i> is a square with edges of length <i>r</i> m, state with a rewhether the centre of mass of the lamina lies within the square or the quarter-circle.	



A rough horizontal rod AB of length 0.45 m rotates with constant angular velocity 6 rad s⁻¹ about a vertical axis through A. A small ring R of mass 0.2 kg can slide on the rod. A particle P of mass 0.1 kg is attached to the mid-point of a light inextensible string of length 0.6 m. One end of the string is attached to R and the other end of the string is attached to R, with angle $RPB = 60^{\circ}$ (see diagram). R and R move in horizontal circles as the system rotates. R is in limiting equilibrium.

(i)	Show that the tension in the portion PR of the string is 1.66 N, correct to 3 significant figures. [5]

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.