

Cambridge Assessment International Education

Cambridge International Advanced Level

| CANDIDATE NAME | | | |
|-----------------------|------------------------|---------------------|---------------------|
| CENTRE NUMBER | | CANDIDATE NUMBER | |
| MATHEMATICS | | | 9709/72 |
| Paper 7 Probability & | Statistics 2 (S2) | Oc | tober/November 2019 |
| | | | 1 hour 15 minutes |
| Candidates answer on | the Question Paper. | | |
| Additional Materials: | List of Formulae (MF9) | | |
| | | | |

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 50.

This document consists of 13 printed pages and 3 blank pages.



BLANK PAGE

| | average, 1 in 150 components made by a certain machine are faulty. The random values the number of faulty components in a random sample of 500 components. | |
|-------|--|--------|
| (i) | Describe fully the distribution of X . | |
| | | |
| | | |
| | | |
| | | |
| (ii) | State a suitable approximating distribution for X , giving a justification for your choice. | |
| | | |
| | | •••••• |
| | | ••••• |
| | | |
| | | |
| | | |
| iii) | Use your approximating distribution to find the probability that the sample will include 3 faulty components. | |
| (iii) | | |
| iii) | | ••••• |
| iii) | 3 faulty components. | ••••• |
| iii) | 3 faulty components. | ••••• |
| iii) | 3 faulty components. | |

The heights of a certain species of animal have been found to have mean 65.2 cm and standard deviation 7.1 cm. A researcher suspects that animals of this species in a certain region are shorter on average than elsewhere. She takes a large random sample of n animals of this species from this region and finds that their mean height is 63.2 cm. She then carries out an appropriate hypothesis test. (i) She finds that the value of the test statistic z is -2.182, correct to 3 decimal places. (a) Stating a necessary assumption, calculate the value of n. [4]

| (b) | Carry out the hypothesis test at the 4% significance level. | [3] |
|------------|---|-------|
| | | ••••• |
| | | |
| | | |
| | | ••••• |
| | | ••••• |
| | | ••••• |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| Exp | plain why it was necessary to use the Central Limit theorem in carrying out the test. | [1] |
| | | |
| | | |
| | | |
| | | |
| | | |

(ii)

| (| Calculate a 96% confidence interval for μ , giving the end-points correct to 1 decimal place. |
|---|---|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| • | |
| • | |
| • | |
| • | |
| • | |

| | • | | | | |
|---|---|--|---|---|---|
| | | | | | |
| • | ••••• | | ••••• | • | ••••• |
| | | | | • | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | • | | | | • |
| | | | | | |
| •••••• | ••••• | ••••• | •••••• | • | • |
| | | | | | |
| | • | | • | • | • |
| | | | | | |
| ••••• | ••••• | ••••• | | ••••• | ••••• |
| | | | | | |
| | | | | • | • |
| | | | | | |
| • | | | | • | |
| | | | | | |
| Without calcula | nple of 50 bags of ation, state when real found in particular and the state of the | ther this confid | lence interval | will be wider | |
| Without calcula | | ther this confid | lence interval | will be wider | |
| Without calcula | ation, state whe | ther this confid | lence interval | will be wider | |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your | will be wider answer. | |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |
| Without calcula | ation, state whe | ther this confid rt (i). Give a re | lence interval ason for your a | will be wider answer. | or narrower tha |

| 4 A random variable <i>X</i> has probability density function given | 4 | A random | variable X ha | s probability | y density | y function | given | by |
|--|---|----------|-----------------|---------------|-----------|------------|-------|----|
|--|---|----------|-----------------|---------------|-----------|------------|-------|----|

s probability density function given by
$$f(x) = \begin{cases} \frac{1}{2}x & 0 \le x \le a, \\ 0 & \text{otherwise,} \end{cases}$$

where a is a constant.

| (i) | Find a. | [2] |
|------|----------------------------------|--------|
| | | |
| | | |
| | | |
| | | •••••• |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | ••••• |
| | | |
| (ii) | Show that $E(X) = \frac{4}{3}$. | [3] |
| ` ' | \ | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | ••••• |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

The median of X is denoted by m.

| i) Find P(I | C(X) < X < m). | | | | [4] |
|-------------|----------------|-------|-------|--------|------|
| | | | | | |
| •••• | | | | | |
| | | | | | |
| | | | | | |
| ••••• | •••••• | ••••• | ••••• | •••••• | |
| | | | | | |
| ••••• | | | | ••••• | |
| | | | | | |
| | | | | ••••• | |
| | | | | | |
| | | | | ••••• | |
| | | | | | |
| | | | | ••••• | |
| | | | | | |
| | | | | | |
| ••••• | •••••• | ••••• | | •••••• | |
| ••••• | •••••• | ••••• | | ••••• | |

| | Find the probability that the total mass of 10 randomly chosen large boxes of chocolates is leading than 3240 g. |
|---|--|
| | |
| | |
| • | |
| • | |
| • | |
| | |
| • | |
| • | |
| • | |
| | |
| | |
| • | |
| • | |
| • | |
| | |
| • | |
| • | |
| • | |
| | |

| wice the mass of a randomly chosen small box of chocolates. | [: |
|---|--------|
| | |
| | |
| | |
| | ••••• |
| | |
| | |
| | |
| | |
| | |
| | |
| | ••••• |
| | |
| | |
| | |
| | |
| | |
| | •••••• |
| | |
| | |
| | |
| | |
| | |
| | |
| | •••••• |
| | |
| | |
| | ••••• |
| | |
| | |

| nas l test v | number of accidents per month, X , at a factory has a Poisson distribution. In the past the mean been 1.1 accidents per month. Some new machinery is introduced and the management wish to whether the mean has increased. They note the number of accidents in a randomly chosen month carry out a hypothesis test at the 1% significance level. |
|-----------------|--|
| (i) | Show that the critical region for the test is $X \ge 5$. Given that the number of accidents is 6, carry out the test. [6] |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |

Later they carry out a similar test, also at the 1% significance level.

| | xplain the meaning of a Type I error in this context and state the proba | |
|-----|--|--|
| ••• | | |
| ••• | | |
| ••• | | |
| | | |
| ••• | | |
| ••• | | |
| ••• | | |
| ••• | | |
| ••• | | |
| ••• | | |
| ••• | | |
| G | iven that the mean is now 7.0, find the probability of a Type II error. | |
| ••• | | |
| ••• | | |
| | | |
| | | |
| ••• | | |
| | | |
| | | |
| | | |
| | | |
| | | |

Additional Page

| If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown. |
|---|
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.