



Cambridge Assessment International Education

Cambridge International Advanced Subsidiary and Advanced Level

CANDIDATE NAME			
CENTRE NUMBER		CANDIDATE NUMBER	
MATHEMATICS			9709/42
Paper 4 Mechanics	1 (M1)		May/June 2019
			1 hour 15 minutes
Candidates answer of	on the Question Paper.		
Additional Materials:	List of Formulae (MF9)		

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

Where a numerical value for the acceleration due to gravity is needed, use 10 m s⁻².

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.


The total number of marks for this paper is 50.





This document consists of 13 printed pages and 3 blank pages.

BLANK PAGE




Coplanar forces of magnitudes $40 \mathrm{N}, 32 \mathrm{N}, P \mathrm{N}$ and $17 \mathrm{N}$ act at a point in the directions shown in the diagram. The system is in equilibrium. Find the values of P and θ .

							••••••
			•••••				· • • • • • • • • • • • • • • • • • • •
			•••••				• • • • • • • • •
							· · · · · · · · · · · · · · · · · · ·
••••••	••••••	•••••	•••••••	•••••	••••••	•••••	• • • • • • • • •
•••••			•••••	•••••	•••••		• • • • • • • •
•••••	•••••	•••••	••••••	•••••	•••••	•••••	••••••
	•••••		••••••	•••••	•••••	•••••	
••••••	•••••	•••••	••••••	•••••	•••••	•••••	· • • • • • • •
•••••••••••	•••••••	•••••	••••••	•••••	•••••	•••••	
							· • • • • • • • • • • • • • • • • • • •

	a line of gred the work d			the parti-	cic a dist	ance or 2	.5 m up u	ie piane	at a con
•••••			•••••				•••••		
			•••••				•••••		
		•••••							
		•••••							
		•••••					•••••		

ca th	constant resistance to motion of magnitude 350 N acts on a car of mass 1250 kg. The engine of the ar exerts a constant driving force of 1200 N. The car travels along a road inclined at an angle of θ to the horizontal, where $\sin \theta = 0.05$. Find the speed of the car when it has moved 100 m from rest in each of the following cases.
•	The car is moving up the hill.
•	The car is moving down the hill. [7]
•••	
•••	
•••	
•••	
•••	
•••	
•••	
•••	
•••	
•••	
•••	
•••	
•••	

• • • • • • • • • • • • • • • • • • • •
• • • • • • • • • • • • • • • • • • • •
• • • • • • • • • • • • • • • • • • • •
• • • • • • • • • • • • • • • • • • • •



Two particles A and B, of masses $0.4 \, \mathrm{kg}$ and $0.2 \, \mathrm{kg}$ respectively, are connected by a light inextensible string which passes over a fixed smooth pulley. Both A and B are $0.5 \, \mathrm{m}$ above the ground. The particles hang vertically (see diagram). The particles are released from rest. In the subsequent motion B does not reach the pulley and A remains at rest after reaching the ground.

(i)	For the motion before A reaches the ground, show that the magnitude of the acceleration of each particle is $\frac{10}{3}$ m s ⁻² and find the tension in the string. [4]

•••••
 •••••
•••••
 •••••
 ••••••
••••••
•••••

		t an angle working at				$\sin \theta = 0$.05, at a ste	eady speed o	of 12 r
•••••	••••••	••••••	•••••	••••••	•••••	••••••	•••••	•••••	•
•••••	••••••	••••••	••••••	•••••	•••••	••••••	••••••	•••••	••••••
•••••			•••••	•••••	•••••	•••••			
•••••	••••••	••••••	•••••	••••••	••••••	•••••	•••••	•••••	
•••••	•••••	•••••			•••••		•••••	•••••	.
•••••									· · · · · · · · · · · · · · · · · · ·

Particles P and Q leave a fixed point A at the same time and travel in the same straight line. The velocity of P after t seconds is 6t(t-3) m s⁻¹ and the velocity of Q after t seconds is (10-2t) m s⁻¹.

(i)	Sketch, on the same axes, velocity-time graphs for P and Q for $0 \le t \le 5$.	[3]
(ii)	Verify that P and Q meet after 5 seconds.	[4
(ii)	Verify that P and Q meet after 5 seconds.	[4
(ii)	Verify that P and Q meet after 5 seconds.	
(ii)		
(ii)		[4]

(iii)	Find the greatest distance between P and Q for $0 \le t \le 5$. [4]

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.