Candidate surname	tails below	v before ente	ering your candidate information Other names
Pearson Edexcel	Centr	e Number	Candidate Number
Tuesday 12 Ja	anı	lary	2021
Afternoon (Time: 1 hour 30 minu	utes)	Paper R	eference WME02/01
Mathematics			
Mathematics International Advance Mechanics M2	ed Sul	osidiar	y/Advanced Level

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets – use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over 🕨



P60706A ©2021 Pearson Education Ltd. 1/1/1/1/1/



1.	A particle <i>P</i> of mass 1.5 kg is moving with velocity $(4\mathbf{i} + 6\mathbf{j})\mathbf{m}\mathbf{s}^{-1}$ when it receives an impulse of magnitude 15 Ns. Immediately after <i>P</i> receives the impulse, the velocity of <i>P</i> is $v\mathbf{i}\mathbf{m}\mathbf{s}^{-1}$.	Lea ^v blar
	Find the two possible values of v . (7)	

DO NOT WRITE IN THIS AREA

Question 1 continued	Leave blank
(Total 7 marks)	Q1

DO NOT WRITE IN THIS AREA

Leave blank

(5)

DO NOT WRITE IN THIS AREA





The uniform lamina ABC has sides AB = AC = 13a and BC = 10a. The lamina is freely suspended from A. A horizontal force of magnitude F is applied to the lamina at B, as shown in Figure 1. The line of action of the force lies in the vertical plane containing the lamina. The lamina is in equilibrium with AB vertical. The weight of the lamina is W.

Find F in terms of W.

2.

Question 2 continued		1
guestion 2 continued		
		Q
		ř
	(Total 5 marks)	

DO NOT WRITE IN THIS AREA

3. A car of mass 600 kg travels along a straight horizontal road with the engine of the car working at a constant rate of *P* watts. The resistance to the motion of the car is modelled as a constant force of magnitude *R* newtons. At the instant when the speed of the car is 15 ms^{-1} , the magnitude of the acceleration of the car is 0.2 ms^{-2} .

Later the same car travels up a straight road inclined at angle θ to the horizontal, where

 $\sin \theta = \frac{1}{20}$. The resistance to the motion of the car from non-gravitational forces is

modelled as a constant force of magnitude *R* newtons. When the engine of the car is working at a constant rate of *P* watts, the car has a constant speed of 10 m s^{-1} .

Find the value of *P*.

(8)

Leave blank

DO NOT WRITE IN THIS AREA

	Leave blank	
Question 3 continued		
		DO
		DO NOT WRITE IN THIS AREA
		WRI
		I
		ARE
		Å
		C
		NC.
		DO NOT WRITE IN THIS ARE
		C
		DO NOT WRITE IN THIS AREA
8		

Question 3 continued		Leave blank
	(Total 8 marks)	Q3
		9 Turn over

Turn over

Leave blank

DO NOT WRITE IN THIS AREA



Figure 2

The number "4", shown in Figure 2, is a rigid framework made from three uniform rods, *AB*, *BC* and *CD*, where

AB = 6a, BC = 5a and CD = 4a

The point *E* is on *AB* and *CD*, where BE = 4a, CE = 3a and angle $CEB = 90^{\circ}$

The three rods are all made from the same material and they all lie in the same plane.

The framework is suspended from *B* and hangs in equilibrium with *BA* at an angle θ to the downward vertical.

Find θ to the nearest degree.

(9)

4.

Question 4 continued	Le bl

DO NOT WRITE IN THIS AREA

lestion 4 continued	
	D
	DO NOT WRITE IN THIS AREA
	WRIT
	E I
	A SIH.
	REA
	DON
	DO NOT WRITE IN THIS ARE
	RITE
	SARE
	Ū
	WR
	DO NOT WRITE IN THIS AREA
	AREA

Question 4 continued		blank
		Q4
	(Total 0 marks)	
	(Total 9 marks)	
		13
		Turn over

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave

At time <i>t</i> seconds, $t \ge 0$, a particle <i>P</i> has velocity $\mathbf{v} \mathbf{m} \mathbf{s}^{-1}$, where	
$\mathbf{v} = (5t^2 - 12t + 15)\mathbf{i} + (t^2 + 8t - 10)\mathbf{j}$	
When $t = 0$, <i>P</i> is at the origin <i>O</i> .	
At time T seconds, P is moving in the direction of $(i + j)$.	
(a) Find the value of <i>T</i> .	(3)
When $t = 3$, P is at the point A.	
(b) Find the magnitude of the acceleration of P as it passes through A .	(4)
(c) Find the position vector of <i>A</i> .	(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

P 6 0 7 0 6 A 0 1 4 2 8

Question 5 continued	Leav blan
(Total 11	

DO NOT WRITE IN THIS AREA

Leave blank

DO NOT WRITE IN THIS AREA



A ladder AB has length 6 m and mass 30 kg. The ladder rests in equilibrium at 60° to the horizontal with the end A on rough horizontal ground and the end B against a smooth vertical wall, as shown in Figure 3.

A man of mass 70 kg stands on the ladder at the point C, where AC = 2 m, and the ladder remains in equilibrium. The ladder is modelled as a uniform rod in a vertical plane perpendicular to the wall. The man is modelled as a particle.

(a) Find the magnitude of the force exerted on the ladder by the ground.

The man climbs further up the ladder. When he is at the point D on the ladder, the ladder is about to slip.

Given that the coefficient of friction between the ladder and the ground is 0.4

(b) find the distance *AD*.

(c) State how you have used the modelling assumption that the ladder is a rod.

(1)

(4)

(6)



Question 6 continued	

DO NOT WRITE IN THIS AREA

unation (continued	Leave blank
uestion 6 continued	
3	

Question 6 continued		Leav
		Q
	(Total 11 marks)	

DO NOT WRITE IN THIS AREA

Leave blank

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





The fixed point *A* is 20 m vertically above the point *O* which is on horizontal ground. At time t = 0, a particle *P* is projected from *A* with speed 12 m s^{-1} at an angle θ° above the horizontal. The particle moves freely under gravity. At time t = 5 seconds, *P* strikes the ground at the point *B*, where OB = 40 m, as shown in Figure 4.

(a) By considering energy, find the speed of P as it hits the ground at B.

(b) Find the least speed of P as it moves from A to B.

(c) Find the length of time for which the speed of P is more than 10 m s^{-1} .

(6)

(4)

(2)

7.

Question 7 continued		

DO NOT WRITE IN THIS AREA

Question 7 continue	d	Leave blank	
22			ノ §

P 6 0 7 0 6 A 0 2 2 2 8

Question 7 continued		
	(Total 12 marks)	r

DO NOT WRITE IN THIS AREA

Leave blank

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

8. Two particles, A and B, have masses 3m and 4m respectively. The particles are moving towards each other along the same straight line on a smooth horizontal surface. The particles collide directly. Immediately after the collision, A and B are moving in the same

direction with speeds $\frac{u}{3}$ and *u* respectively. In the collision, *A* receives an impulse of magnitude 8*mu*.

(a) Find the coefficient of restitution between A and B.

(6)

(6)

When A and B collide they are at a distance d from a smooth vertical wall, which is perpendicular to their direction of motion. After the collision with A, particle B collides directly with the wall and rebounds so that there is a second collision between A and B. This second collision takes place at distance x from the wall.

Given that the coefficient of restitution between B and the wall is $\frac{1}{4}$

(b) find x in terms of d.

Question 8 continued	Lea bla

DO NOT WRITE IN THIS AREA

Question 8 continued	Leav blan	ve Ik

Question 8 continued	

DO NOT WRITE IN THIS AREA

Question 8 continued	
(Total 12 marks)	00 1
TOTAL FOR PAPER: 75 MARKS	Q8
END	