

Cambridge O Level

	CANDIDATE NAME						
	CENTRE NUMBER		CANDIDATE NUMBER				
* ω Ν Φ Φ Φ Φ Φ Φ Φ Φ Φ Φ Φ Φ Φ Φ Φ Φ Φ Φ	CHEMISTRY			5070/41			
ω	Paper 4 Alternat	tive to Practical	Oct	October/November 2020			
0 0				1 hour			
о 7 N 0 0 0 0	You must answe	er on the question paper.					
٥	No additional m	paterials are needed					

No additional materials are needed.

INSTRUCTIONS

- Answer all questions. •
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs. •
- Write your name, centre number and candidate number in the boxes at the top of the page. •
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid. •
- Do not write on any bar codes. •
- You may use a calculator. •
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 60.
- The number of marks for each question or part question is shown in brackets [].

1 A student wants to separate a mixture of dyes so that the dyes in the mixture can be identified.

The apparatus the student uses is shown.

The student also tests dyes **A**, **B**, **C** and **D**.



(a) Name the process the student uses to separate the mixture of dyes.

(b) The apparatus shown in the diagram is set up incorrectly. Explain why.

......[1]

The apparatus is then set up correctly.

The solvent travels up the paper.

The paper is removed and then dried.

The result is shown in the diagram.



(c) Which of the dyes A, B, C and D are present in the mixture of dyes?

(ii) Calculate the $R_{\rm f}$ value of **A**.

[Total: 5]

[1]

2 When metal carbonates are heated they decompose and give off a gas that turns limewater milky.

A student investigates the rate of decomposition of four metal carbonates using the apparatus shown.



The student:

- separately heats each of the metal carbonates with a Bunsen burner
- measures the time taken for the limewater to turn milky.

The results are shown in the table.

metal carbonate	time taken for limewater to turn milky/s
calcium carbonate	500
lead(II) carbonate	120
magnesium carbonate	300
zinc carbonate	205

(a) Name the gas that turns limewater milky.

......[1]

(b)	The	The amount of heat supplied by the Bunsen burner must be controlled in each experiment.					
	(i)	Suggest two ways in which the heat supplied by the Bunsen burner is kept constant.					
		1					
		2[2]					
	(ii)	Suggest two other variables that must be controlled.					
		1					
		2[2]					
(c)	Wh	ich metal carbonate is the slowest to decompose?					
		[1]					
(d)	Sug	gest how the investigation can be improved to make the results more reliable.					
		[2]					
		[Total: 8]					

3 A student is provided with dilute sulfuric acid labelled L.

The student determines the concentration of L by titration.

The student uses apparatus **A** to measure and transfer 25.0 cm^3 of **L** into apparatus **B**.

The diagrams show apparatus **A** and **B**.



Solution **R** is $0.100 \text{ mol}/\text{dm}^3$ aqueous sodium hydroxide, NaOH(aq).

R is put into a burette and run into the conical flask until the indicator changes colour and the end-point is reached.

- (c) Which liquid is used to wash out the burette immediately before it is filled with R?
 -[1]
- (d) What is the colour change of the methyl orange indicator at the end-point?

The colour changes from [1]

(e) The student does three titrations. The diagrams show parts of the burette with the liquid levels at the beginning and end of each titration.



(i) Use the diagrams to complete the table.

titration number	1	2	3
final burette reading/cm ³			
initial burette reading/cm ³			
volume of R /cm ³			
best titration results (\checkmark)			

Summary

Tick (\checkmark) the best titration results in the table.

Use the ticked values to calculate the average volume of R.

..... cm³ [4]

(ii) Why was it unnecessary for the student to do a third titration?

(f) Solution **R** is 0.100 mol/dm³ aqueous sodium hydroxide, NaOH(aq).

Calculate the number of moles of NaOH in the average volume of 0.100 mol/dm³NaOH used in the titration. Give your answer to **three** significant figures.

..... mol [1]

(g) Dilute sulfuric acid reacts with aqueous sodium hydroxide as shown.

$$H_2SO_4 + 2NaOH \rightarrow Na_2SO_4 + 2H_2O$$

Calculate the number of moles of H_2SO_4 in 25.0 cm³ of **M**.

..... mol [1]

(h) Calculate the number of moles of H_2SO_4 in 500 cm³ of **M**.

..... mol [1]

(i) Calculate the number of moles of H_2SO_4 in 25.0 cm³ of L.

..... mol [1]

(j) Calculate the concentration of H_2SO_4 in L.

..... mol/dm³ [1]

- (k) Between each titration the student washes the conical flask with water and dries it.
 - (i) A second student repeats the experiment but uses L instead of water to wash the conical flask and does not dry the flask.

State **and** explain whether the second student's titration volume of NaOH(aq) would be smaller, larger or unchanged compared with the first student's titration volume of NaOH(aq).

(ii) A third student repeats the experiment, washes the conical flask out with water but does not dry it before each titration.

State **and** explain whether the third student's titration volume of NaOH(aq) would be smaller, larger or unchanged compared with the first student's titration volume of NaOH(aq).

|
 |
|------|------|------|------|------|------|------|---------|
|
 |
|
 |
[2] |

[Total: 20]

4 A student wants to electroplate a metallic spoon with a layer of copper.

Plan how the student does the experiment.

You should:

- state what needs to be done to the spoon before electrolysis
- include a labelled diagram of the apparatus
- give the names of the substances used.

 	[5]

5 Compound **Q** contains two cations and one anion. The following table shows the tests a student does on compound **Q**.

Complete the table.

Identify any gases that are formed in the tests.

	tes	st	observation	conclusion	
 (a) Q is dissolved in water. The solution is divided into two portions for tests (b) and (c). 		n is divided rtions for tests	A coloured solution is formed.		
					[1]
(b)	aqueo hydrox	first portion, us sodium kide is added		Q contains Cr ³⁺ or Fe ²⁺ ions.	
	until a seen.	change is			[1]
	hydrox	us sodium kide is added		Q contains Fe ²⁺ ions.	
	to the (b)(i) .	mixture from			[1]
		ixture from is warmed and s formed is	The gas turns damp red litmus paper blue.		-
	tested litmus	with damp red paper.			
					[2]
(c)				Q contains SO_4^{2-} ions.	
					[3]

[Total: 8]

The diagram shows the apparatus the student uses.



(a) The student burns a known mass of ethanol.

The heat released is used to heat 200 cm³ of water.

The student determines the temperature rise of the water while the ethanol is burning.

(i) In the first experiment the student burns 1.0g of ethanol and the temperature rises by 20 °C.

State all the measurements the student makes to obtain these values.

- (ii) Why is a metal beaker used instead of a glass beaker?
 -[1]
- (iii) The student finds out that 1.0g of ethanol should give a theoretical temperature rise of more than 20 °C.

Suggest **two** reasons why the temperature rise of the water was less than the theoretical value.

1 2 [2]

(iv) Suggest two ways, using 1.0 g of ethanol and 200 cm³ of water, in which the apparatus can be changed to produce a temperature rise of more than 20 °C.

1 2 [2]

(b) The relative formula mass of ethanol, C_2H_5OH , is 46.

Calculate the number of moles of ethanol in 1.0 g.

..... mol [1]

14

(c) The experiment is repeated using the same number of moles of different alcohols.

The temperature rises are shown in the table.

number of carbon atoms in each alcohol molecule	temperature rise/°C
2	20
3	32
4	43
5	55
6	66
7	78

(i) What evidence in the table shows that all the reactions are exothermic?

......[1]

(ii) Plot the data in the table on the grid.

Draw a straight line of best fit.

Extend your line to the left hand side of the grid.



(d) The student repeats the experiment with methanol, CH_3OH .

The student uses the same number of moles of methanol as the other alcohols and the same volume of water.

Use your graph to predict the temperature rise for methanol.

.....°C [1]

- (e) The student predicts that the temperature rise with another alcohol is 90 °C.
 - (i) The student uses an initial water temperature of 20 °C.

Explain why it is not possible to obtain a temperature rise of 90 °C.

......[1]

(ii) The student wants to obtain a temperature rise of 90 °C with this alcohol and the same apparatus.

Suggest one way the student can obtain a temperature rise of 90 °C.

[Total: 14]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.